arXiv:math/0604592v1 [math.AT] 27 Apr 2006

A CUBICAL APPROACH TO HOMOTOPY ORBITS OF CIRCLE ACTIONS

Kathryn Hess

Ecole Polytechnique Fédérale de Lausanne

27.04.06

Abstract. For a space X belonging to a class including all double suspensions of finite type, we construct for any field $\Bbbk$ a cochain complex $hos^*(X)$ such that $\mathrm{H}^*\big(hos^*(X)\big) \cong \mathrm{H}^*(\mathcal{L}X_{hS^1};\Bbbk)$, the cohomology of the homotopy orbit space of the natural S^1-action on the free loop space $\mathcal{L}X$. The model $hos^*(X)$ is "small" and thus lends itself relatively easily to explicit computation. Furthermore $hos^*(X)$ fits into an extension of cochain complexes that models the fibration $\mathcal{L}X \longrightarrow \mathcal{L}X_{hS^1} \longrightarrow BS^1$. This concrete link with topology proves essential for the construction of the cochain complex for computing spectrum cohomology of topological cyclic homology in [HR].

Introduction

For a space X belonging to a class including all double suspensions of finite type, we construct here for any field $\Bbbk$ a cochain complex $hos^*(X)$ such that $\mathrm{H}^*\big(hos^*(X)\big) \cong \mathrm{H}^*(\mathcal{L}X_{hS^1};\Bbbk)$, the cohomology of the homotopy orbit space of the natural S^1-action on the free loop space $\mathcal{L}X$, also known as the equivariant or Borel cohomology of the free loop space. The model $hos^*(X)$ is "small" and thus lends itself relatively easily to explicit computation, as we illustrate in section 2.4. Furthermore, $hos^*(X)$ fits into a commuting diagram

(A)
$$\begin{array}{ccccc}
\Bbbk[\upsilon] & \longrightarrow & hos^*(X) & \longrightarrow & fls^*(X) \\
\downarrow{\scriptstyle =} & & \downarrow{\scriptstyle \simeq} & & \downarrow{\scriptstyle \simeq} \\
\Bbbk[\upsilon] & \longrightarrow & \bullet & \longrightarrow & CU^*(\mathcal{L}X,\Bbbk) \\
\uparrow{\scriptstyle \simeq} & & \uparrow{\scriptstyle \simeq} & & \uparrow{\scriptstyle =} \\
CU^*(BS^1;\Bbbk) & \longrightarrow & CU^*(\mathcal{L}X_{hS^1};\Bbbk) & \xrightarrow{CU^*i} & CU^*(\mathcal{L}X,\Bbbk),
\end{array}$$

where $fls^*(X)$ is a simplified version of the free loop space model of [DH] and [H], $\Bbbk[\upsilon]$ is a polynomial algebra on a generator of degree 2, CU^* denotes cubical cochains and $\mathcal{L}X \longrightarrow \mathcal{L}X_{hS^1} \longrightarrow BS^1$ is the usual fibration sequence. The

1991 *Mathematics Subject Classification.* Primary: 55R20 Secondary: 18D50, 18G30, 18G35, 18G55, 55P35, 55P40, 55U05, 55U10, 57T05.

Key words and phrases. Circle action, homotopy orbits, free loop space, cubical chains, strongly homotopy morphism, operad.

Work completed during a visit to the Institut Mittag-Leffler (Djursholm, Sweden).

existence of this concrete link with topology proves essential for the construction of the cochain complex for computing spectrum cohomology of topological cyclic homology in [HR].

We begin in the first section of this paper with a general discussion of the construction of complexes for calculation of equivariant (co)homology. After recalling the operadic description of "strongly homotopy" structures from [HPS1] and [HPST], we introduce a more highly structured notion of resolution, which we call DCSH resolution, leading to an enriched version of Moore's famous theorem (Theorem 1.1.2). In particular, we show that if G is a connected topological group, E is the total space of a principal G-bundle and Y is a G-space, then we can calculate the algebra structure of $\mathrm{H}^*(E \underset{G}{\times} Y)$, given DCSH resolutions of CU_*G and of the cubical chains on CU_*E as a CU_*G-module.

After proving the existence of a very special family of primitive elements in the reduced cubical chains on S^1 and studying its properties, we introduce a particularly useful DCSH resolution of the cubical chains on ES^1 as a module over the cubical chains on S^1 (1.3.1). Applying the general construction of our enriched Moore's Theorem and dualizing, we obtain a cochain complex for computing the graded algebra $\mathrm{H}^* Y_{hS^1}$ for any S^1-space Y (1.4.3). The Batalin-Vilkovisky structure on $\mathrm{H}^* Y$ shows up in this complex, as one summand of the differential.

In section 2, we specialize to the case of $\mathcal{L}X$, obtaining a small cochain model for $\mathcal{L}X_{hS^1}$ as an extension of a model of the free loop space. The first part of section 2 is devoted to refining and simplifying the free loop model of [DH] and [H], leading to the small, "linearized" model $fls^*(X)$ of Theorem 2.2.1. Blending the methods of section 1 and the model $fls^*(X)$, we then obtain the desired model $hos^*(X)$ (Theorem 2.3.3).

In proving the existence of the commuting diagram (A), we apply an apparently new result from homological perturbation theory (Lemma 2.3.4) that we think may be useful in other contexts as well. Indeed it has already played a crucial role in [HR], where it is applied to constructing a model of the p^{th}-power map on the free loop space.

Throughout this paper, we elaborate upon and improve results from [H]. In particular, the operadic perspective on computing algebra structure of equivariant cohomology is new.

The author is deeply grateful to Nicolas Dupont for the ideas he contributed during our discussions of $\mathcal{L}X_{hS^1}$ during the 1990's.

0. Preliminaries

In this section we first recall definitions and constructions and fix our notation and terminology. We then outline the operadic approach to the study of strongly homotopy structures, as developed in [HPS] and [HPST].

0.1 Elementary definitions, terminology and notation.

Given a category $\mathbf{C}$ and objects A and B, the class of morphisms from A to B is denoted $\mathbf{C}(A, B)$.

In any model category we use the notation $\rightarrowtail$ for cofibrations, $\twoheadrightarrow$ for fibrations and $\xrightarrow{\sim}$ for weak equivalences.

Chain and cochain complexes. We write $CU_*(X)$ to denote the reduced cubical chains on a topological space X.

Throughout this paper we work over a principal ideal domain R. Dualization is indicated by a $\sharp$ as superscript. The degree of an element x in a graded module is denoted $|x|$, unless it is used as an exponent, in which case the bars are dropped, if there is no confusion possible.

The Koszul sign convention for commuting elements of a graded module or for commuting a morphism of graded modules past an element of the source module is used consistently throughout this article. For example, if V and W are graded algebras and $v \otimes w, v' \otimes w' \in V \otimes W$, then

$$(v \otimes w) \cdot (v' \otimes w') = (-1)^{|w|\cdot|v'|} vv' \otimes ww'.$$

Futhermore, if $f : V \to V'$ and $g : W \to W'$ are morphisms of graded modules, then for all $v \otimes w \in V \otimes W$,

$$(f \otimes g)(v \otimes w) = (-1)^{|g|\cdot|v|} f(v) \otimes g(w).$$

The source of the Koszul sign convention is the definition of the twisting isomorphism

$$\tau : V \otimes W \to W \otimes V : v \otimes w \mapsto (-1)^{|v|\cdot|w|} w \otimes v.$$

Given chain complexes (V, d) and (W, d), the notation $f : (V, d) \xrightarrow{\simeq} (W, d)$ indicates that f induces an isomorphism in homology. In this case we refer to f as a *quasi-isomorphism*.

If $V = \bigoplus_{i\in\mathbb{Z}} V_i$ is a graded module, then $s^{-1}V$ and sV denote the graded modules with, respectively, $(s^{-1}V)_i \cong V_{i+1}$ and $(sV)_i \cong V_{i-1}$. Given a homogeneous element v in V, we write $s^{-1}v$ and sv for the corresponding elements of $s^{-1}V$ and sV. If the gradings are written as upper indices, i.e., $V = \bigoplus_{i\in\mathbb{Z}} V^i$, then $(s^{-1}V)^i \cong V^{i-1}$ and $(sV)^i \cong V^{i+1}$.

A graded R-module $V = \bigoplus_{i\in\mathbb{Z}} V_i$ is *connected* if $V_{<0} = 0$ and $V_0 \cong R$. It is *simply connected* if, in addition, $V_1 = 0$. We write V_+ for $V_{>0}$. It is of *finite type* if V_i is finitely generated as an R-module for all i.

Chain and cochain algebras and coalgebras over R. Let V be a positively graded, free R-module. The free associative algebra on V is denoted TV, i.e.,

$$TV \cong R \oplus V \oplus (V \otimes V) \oplus (V \otimes V \otimes V) \oplus \cdots.$$

A typical basis element of TV is denoted $v_1 \cdots v_n$, i.e., we drop the tensors from the notation. The product on TV is then defined by

$$\mu(u_1 \cdots u_m \otimes v_1 \cdots v_n) = u_1 \cdots u_m v_1 \cdots v_n.$$

The cofree, coassociative coalgebra on V, denoted $\perp V$ in this article, is isomorphic as a graded R-module to TV. We write $\perp^n V = \bigotimes^n V$, of which a typical basis element is denoted $v_1|\cdots|v_n$. The coproduct on $\perp V$ is then defined in the obvious manner by

$$\begin{aligned}\Delta(v_1|\cdots|v_n) = {} & v_1|\cdots|v_n \otimes 1 + 1 \otimes v_1|\cdots|v_n \\ & + \sum_{i=1}^{n-1} v_1|\cdots|v_i \otimes v_{i+1}|\cdots|v_n.\end{aligned}$$

Let (C,d) be a simply connected (co)chain coalgebra with reduced coproduct $\overline{\Delta}$. The *cobar construction* on (C,d), denoted $\Omega(C,d)$, is the (co)chain algebra $(Ts^{-1}(C_+),d_\Omega)$, where $d_\Omega=-s^{-1}ds+(s^{-1}\otimes s^{-1})\overline{\Delta}s$ on generators. The cobar construction defines a functor from the category of simply connected (co)chain coalgebras to the category of connected (co)chain algebras.

As Milgram showed in [Mi], for every pair of simply connected chain coalgebras (C,d) and (C',d'), there is a quasi-isomorphism of chain algebras

$$q:\Omega\big((C,d)\otimes(C',d')\big)\longrightarrow\Omega(C,d)\otimes\Omega(C',d')$$

specified by $q\big(s^{-1}(x\otimes 1)\big)=s^{-1}x\otimes 1$, $q\big(s^{-1}(1\otimes y)\big)=1\otimes s^{-1}y$ and $q\big(s^{-1}(x\otimes y)\big)=0$ for all $x\in C_+$ and $y\in C'_+$.

Let (A,d) be a connected chain algebra or a simply connected cochain algebra over R, and let $\bar{A}$ be the component of A of positive degree. The *bar construction* on (A,d), denoted $\mathcal{B}(A,d)$, is a differential graded coalgebra $(\perp(s\bar{A}),D_{\mathcal{B}})$. Let $(D_{\mathcal{B}})_1$ denote the linear part of the differential, i.e., $(D_{\mathcal{B}})_1=\pi D_{\mathcal{B}}$, where $\pi:\perp V\longrightarrow V$ is the natural projection. The linear part of $D_{\mathcal{B}}$ specifies the entire differential and is given by

$$(D_{\mathcal{B}})_1(sa_1|\cdots|sa_n)=\begin{cases}-s(da_1) & \text{if}\quad n=1\\ (-1)^{a_1+1}s(a_1\cdot a_2) & \text{if}\quad n=2\\ 0 & \text{if}\quad n>2.\end{cases}$$

The bar and cobar constructions form an adjoint pair of functors, where the bar construction is the right adjoint. We denote the counit of the adjunction by $\varepsilon:\Omega\mathcal{B}\longrightarrow Id$.

Definition. Let A be a chain algebra. A chain complex M is a (right) *A-module coalgebra* if M admits a chain coalgebra structure and a (right) A-module structure such that the A-action map $M\otimes A\longrightarrow A$ is a coalgebra map.

Definition. Let A be a chain algebra. A right A-module M is *semifree* if it is the union of an increasing family $M(-1)=0\subseteq M(0)\subseteq M(1)\subseteq\ldots$ such that each quotient $M(n)/M(n-1)$ is A-free on a basis of cycles.

In particular, if M is semifree, then there is a graded module V such that $M\cong V\otimes A$ as graded modules. There is an analogous notion of semifree left A-modules. We refer the reader to [FHT]for further details.

Throughout section 2 of this article we make extensive use of the following notion.

Definition. Let (A,d) be a cochain R-algebra, and let (B,d) be a cochain complex. A *twisted algebra extension of (A,d) by (B,d)* is a cochain R-algebra $(A\odot B,D)$ such that

(1) the graded R-module underlying $A\odot B$ is $A\otimes B$;
(2) (A,d) is a cochain subalgebra of $(A\odot B,D)$ with respect to the natural inclusion $A\hookrightarrow A\odot B$, i.e., $D(a\otimes 1)=da\otimes 1$ and $(a\otimes 1)\cdot(a'\otimes 1)=aa'\otimes 1$ for all $a,a'\in A$;
(3) (B,d) is a quotient cochain algebra of $(A\odot B,D)$ with respect to the natural projection from $A\odot B$, i.e., $D(1\otimes b)-1\otimes db\in A^+\otimes B$ and $(1\otimes b)\cdot(1\otimes b')-1\otimes bb'\in A^+\otimes B$;
(4) the right action of A on $A\odot B$ satisfies $(1\otimes b)\cdot(a\otimes 1)=(-1)^{|a|\cdot|b|}a\otimes b$ for all $a\in A$, $b\in B$;

(5) the left action of A on $A \odot B$ commutes with the right action and satisfies $(a \otimes 1) \cdot (1 \otimes b) - a \otimes b \in A^{>|a|} \otimes B$ for all $a \in A$, $b \in B$;
(6) D commutes with the right action of A, i.e., $D(a \otimes b) = (-1)^{|a| \cdot |b|}\big(D(1 \otimes b) \cdot (a \otimes 1) + (-1)^b (1 \otimes b) \cdot D(a \otimes 1)$ for all $a \in A$, $b \in B$.

A linear map $f : (A, d) \to (B, d)$, where (A, d) and (B, d) are (co)chain algebras, is a *quasi-algebra map* if f induces a map of algebras in (co)homology.

Simplicial sets. We say that a simplicial set is of *finite type* if it has a finite number of nondegenerate simplices in each dimension.

Definition. Let K be a simplicial set, and let $\mathcal{F}_{ab}$ denote the free abelian group functor. For all $n > 0$, let $DK_n = \cup_{i=0}^{n-1} s_i(K_{n-1})$, the set of degenerate n-simplices of K. The *normalized chain complex* on K, denoted $C_*(K)$, is given by

$$C_n(K) = \mathcal{F}_{ab}(K_n)/\mathcal{F}_{ab}(DK_n).$$

Given a map of simplicial sets $f : K \to L$, the induced map of normalized chain complexes is denoted $C_* f$.

Definition. Let K be a reduced simplicial set, and let $\mathcal{F}$ denote the free group functor. The *loop group* GK on K is the simplicial group such that $(GK)_n = \mathcal{F}(K_{n+1} \setminus Ims_0)$, with faces and degeneracies specified by

$$\begin{aligned} \partial_0 \bar{x} &= (\ \overline{\partial_0 x}\)^{-1} \overline{\partial_1 x} \\ \partial_i \bar{x} &= \overline{\partial_{i+1} x} \quad \text{for all } i > 0 \\ s_i \bar{x} &= \overline{s_{i+1} x} \quad \text{for all } i \geq 0 \end{aligned}$$

where $\bar{x}$ denotes the class in $(GK)_n$ of $x \in K_{n+1}$.

We write $|K|$ for the *geometric realization* of a simplicial set K and $\mathcal{S}_\bullet X$ for the simplicial set of singular simplices of a topological space X. Recall that $(|-|, \mathcal{S}_\bullet)$ is an adjoint pair of functors such that the unit and counit are weak equivalences. We let $S_* X := C_* \circ \mathcal{S}_\bullet X$.

0.2 The category DCSH.

The category **DCSH** of coassociative chain coalgebras and of coalgebra morphisms up to strong homotopy was first defined by Gugenheim and Munkholm in the early 1970's [GM], when they were studying extended naturality of the functor Cotor. The objects of **DCSH** have a relatively simple algebraic description, while that of the morphisms is rich and complex. Its objects are augmented, coassociative chain coalgebras, and a morphism from C to C' is a map of chain algebras $\Omega C \to \Omega C'$.

In a slight abuse of terminology, we say that a chain map between chain coalgebras $f : C \to C'$ is a *DCSH map* if there is a morphism in $\mathbf{DCSH}(C, C')$ of which f is the linear part. In other words, there is a map of chain algebras $g : \Omega C \to \Omega C'$ such that

$$g|_{s^{-1}C_+} = s^{-1} f s + \text{higher-order terms.}$$

In a further abuse of notation, we sometimes write $\widetilde{\Omega} f : \Omega C \to \Omega C'$ to indicate one choice of chain algebra map of which f is the linear part.

The category **DCSH** plays an important role in topology. As established in [HPST], for any reduced simplicial set K, the usual coproduct on $C(K)$ is a DCSH map. Furthermore, given any natural, strictly coassociative coproduct on $\Omega C(K)$, any natural map of chain algebras $\Omega C(K) \to C(GK)$ is also a DCSH map.

In [HPS1] the authors provided a purely operadic description of **DCSH**. Before recalling this description, we briefly explain the framework in which it is constructed. We refer the reader to Section 2 of [HPS1] for further detailes.

Let $\mathbf{M}$ denote the category of chain complexes over a PID R, and let $\mathbf{M}^\Sigma$denote the category of symmetric sequences of chain complexes. An object $\mathcal{X}$ of $\mathbf{M}^\Sigma$ is a family $\{\mathcal{X}(n) \in \mathbf{M} \mid n \geq 0\}$ of objects in $\mathbf{M}$ such that $\mathcal{X}(n)$ admits a right action of the symmetric group Σ_n, for all n. The object $\mathcal{X}(n)$ is called the n^{th}-*level* of the symmetric sequence $\mathcal{X}$. Given a morphism of symmetric sequences $\varphi : \mathcal{X} \to \mathcal{Y}$, we let $\varphi(n) : \mathcal{X}(n) \to \mathcal{Y}(n)$ denote its restriction to level n.

There is a faithful functor $\mathcal{T} : \mathbf{M} \longrightarrow \mathbf{M}^\Sigma$ where, for all n, $\mathcal{T}(A)(n) = A^{\otimes n}$, where Σ_n acts by permuting the tensor factors. The functor $\mathcal{T}$ is strong monoidal, with respect to the *level monoidal structure* $(\mathbf{M}^\Sigma, \otimes, \mathcal{C})$, where $(\mathcal{X} \otimes \mathcal{Y})(n) = \mathcal{X}(n) \otimes \mathcal{Y}(n)$, endowed with the diagonal action of Σ_n, and $\mathcal{C}(n) = R$, endowed with the trivial Σ_n-action.

The category $\mathbf{M}^\Sigma$ also admits a nonsymmetric, right-closed monoidal structure $(\mathbf{M}^\Sigma, \diamond, \mathcal{J})$, where $\diamond$ is the *composition product* of symmetric sequences, and $\mathcal{J}(1) = R$ and $\mathcal{J}(n) = 0$ otherwise. Given symmetric sequences $\mathcal{X}$ and $\mathcal{Y}$,

$$(\mathcal{X} \diamond \mathcal{Y})(n) = \coprod_{\substack{k \geq 1 \\ \vec{n} \in I_{k,n}}} \mathcal{X}(k) \underset{\Sigma_k}{\otimes} \big(Y(n_1) \otimes \cdots \otimes Y(n_k)\big) \underset{\Sigma_{\vec{\imath}}}{\otimes} R[\Sigma_n],$$

where $I_{k,n} = \{\vec{n} = (n_1, ..., n_k) \in \mathbb{N}^k \mid \sum_j n_j = n, n_j \geq 0 \, \forall j\}$ and $\Sigma_{\vec{n}} = \Sigma_{n_1} \times \cdots \times \Sigma_{n_k}$, seen as a subgroup of Σ_n. For any objects $\mathcal{X}, \mathcal{X}', \mathcal{Y}, \mathcal{Y}'$ in $\mathbf{M}^\Sigma$, there is an obvious, natural intertwining map

$$\iota : (\mathcal{X} \otimes \mathcal{X}') \diamond (\mathcal{Y} \otimes \mathcal{Y}') \longrightarrow (\mathcal{X} \diamond \mathcal{Y}) \otimes (\mathcal{X}' \diamond \mathcal{Y}') \, . \tag{0.2.1}$$

An *operad* in $\mathbf{M}$ is a monoid with respect to the composition product. The *associative operad* $\mathcal{A}$ is given by $\mathcal{A}(n) = R[\Sigma_n]$ for all n, endowed with the obvious monoidal structure, induced by permutation of blocks.

Let $\mathcal{P}$ denote any operad in $\mathbf{M}$. A $\mathcal{P}$-*coalgebra* consists of an object C in $\mathbf{M}$. together with an appropriately equivariant and associative family

$$\{C \otimes \mathcal{P}(n) \longrightarrow C^{\otimes n} \mid n \geq 0\}$$

of morphisms in $\mathbf{M}$. The functor $\mathcal{T}$ restricts to a faithful functor

$$\mathcal{T} : \mathcal{P}\text{-}\mathbf{Coalg} \longrightarrow \mathbf{Mod}_{\mathcal{P}}$$

from the category of $\mathcal{P}$-coalgebras to the category of right $\mathcal{P}$-modules.

Given a right $\mathcal{P}$-module $\mathcal{M}$ and a left $\mathcal{P}$-module $\mathcal{N}$, we can define their *composition product over* $\mathcal{P}$, denoted $\mathcal{M} \underset{\mathcal{P}}{\diamond} \mathcal{N}$, to be the coequalizer of the two obvious maps $\mathcal{M} \diamond \mathcal{P} \diamond \mathcal{N} \to \mathcal{M} \diamond \mathcal{N}$ induced by the right and left actions of $\mathcal{P}$.

In [HPS1] the authors constructed a semifree $\mathcal{A}$-bimodule $\mathcal{F}$, called the *Alexander-Whitney bimodule*. As symmetric sequences of graded modules, $\mathcal{F} = \mathcal{A} \diamond \mathcal{S} \diamond \mathcal{A}$,

where $\mathcal{S}(n) = R[\Sigma_n] \cdot z_{n-1}$, the free $R[\Sigma_n]$-module on a generator of degree $n-1$. Moreover, $\mathcal{F}$ admits an increasing, differential filtration, given by $F_n\mathcal{F} = \mathcal{A}\diamond\mathcal{S}_n\diamond\mathcal{A}$, where $\mathcal{S}_n(m) = \mathcal{S}(m)$ if $m \leq n$ and $\mathcal{S}_n(m) = 0$ otherwise. More precisely, if $\partial_{\mathcal{F}}$ is the differential on $\mathcal{F}$, then

$$\partial_{\mathcal{F}} z_n = \sum_{0\leq i\leq n-1} \delta \otimes (z_i \otimes z_{n-i-1}) + \sum_{0\leq i\leq n-1} z_{n-1} \otimes (1^{\otimes i} \otimes \delta \otimes 1^{\otimes n-i-1}),$$

where $\delta \in \mathcal{A}(2) = R[\Sigma_2]$ is a generator.

The Alexander-Whitney bimodule is endowed with a coassociative, counital coproduct

$$\psi_{\mathcal{F}} : \mathcal{F} \to \mathcal{F} \underset{\mathcal{A}}{\diamond} \mathcal{F},$$

where $\underset{\mathcal{A}}{\diamond}$ denotes the composition product over $\mathcal{A}$, defined as the obvious coequalizer. In particular,

$$\psi_{\mathcal{F}}(z_n) = \sum_{\substack{1\leq k\leq n+1 \\ \vec{\imath}\in I_{k,n+1}}} z_{k-1} \otimes (z_{i_1-1} \otimes \cdots \otimes z_{i_k-1})$$

for all $n \geq 0$, where $I_{k,n} = \{\vec{\imath} = (i_1, ..., i_k) \mid \sum_j i_j = n\}$.

Furthermore, $\mathcal{F}$ is a level comonoid, i.e., there is a coassociative, counital coproduct

$$\Delta_{\mathcal{F}} : \mathcal{F} \longrightarrow \mathcal{F} \otimes \mathcal{F},$$

which is specified by

$$\Delta_{\mathcal{F}}(z_n) = \sum_{\substack{1\leq k\leq n+1 \\ \vec{\imath}\in I_{k,n+1}}} \big(z_{k-1} \otimes (\delta^{(i_1)} \otimes \cdots \otimes \delta^{(i_k)})\big) \otimes \big(\delta^{(k)} \otimes (z_{i_1-1} \otimes \cdots \otimes z_{i_k-1})\big).$$

Here, $\delta^{(i)} \in \mathcal{A}(i)$ denotes the appropriate iterated composition product of $\delta^{(2)} = \delta$.

Let $(\mathcal{A}, \psi_{\mathcal{F}})$-**Coalg** denote the category of which the objects are $\mathcal{A}$-coalgebras (i.e., coassociative and counital chain coalgebras) and where the morphisms are defined by

$$(\mathcal{A}, \psi_{\mathcal{F}})\text{-}\mathbf{Coalg}(C, C') := \mathbf{Mod}_{\mathcal{A}}\big(\mathcal{T}(C) \underset{\mathcal{A}}{\diamond} \mathcal{F}, \mathcal{T}(C')\big).$$

Composition in $(\mathcal{A}, \psi_{\mathcal{F}})$-**Coalg** is defined in terms of $\psi_{\mathcal{F}}$. Given $\theta \in (\mathcal{A}, \psi_{\mathcal{F}})\text{-}\mathbf{Coalg}(C, C')$ and $\theta' \in (\mathcal{A}, \psi_{\mathcal{F}})\text{-}\mathbf{Coalg}(C', C'')$, their composite $\theta'\theta \in (\mathcal{A}, \psi_{\mathcal{F}})\text{-}\mathbf{Coalg}(C, C'')$ is given by composing the following sequence of (strict) morphisms of right $\mathcal{A}$-modules.

$$\mathcal{T}(C) \underset{\mathcal{A}}{\diamond} \mathcal{F} \xrightarrow{1_{\mathcal{T}(C)} \underset{\mathcal{A}}{\diamond} \psi_{\mathcal{F}}} \mathcal{T}(C) \underset{\mathcal{A}}{\diamond} \mathcal{F} \underset{\mathcal{A}}{\diamond} \mathcal{F} \xrightarrow{\theta \underset{\mathcal{A}}{\diamond} 1_{\mathcal{F}}} \mathcal{T}(C') \underset{\mathcal{A}}{\diamond} \mathcal{F} \xrightarrow{\theta'} \mathcal{T}(C'').$$


We call $(\mathcal{A}, \psi_{\mathcal{F}})$-**Coalg** the *$\psi_{\mathcal{F}}$-governed category of $\mathcal{A}$-coalgebras*.

Theorem 0.2.1 [HPS1]**.** *There is an isomorphism of categories*

$$(\mathcal{A}, \psi)\text{-}\mathbf{Coalg}(C, C') \cong \mathbf{DCSH}.$$

Define a bifunctor $\wedge : (\mathcal{A}, \psi_{\mathcal{F}})\text{-}\mathbf{Coalg} \times (\mathcal{A}, \psi_{\mathcal{F}})\text{-}\mathbf{Coalg} \to (\mathcal{A}, \psi_{\mathcal{F}})\text{-}\mathbf{Coalg}$ on objects by $C \wedge C' := C \otimes C'$, the usual tensor product of chain coalgebras. Given

$\theta \in (\mathcal{A},\psi_{\mathcal{F}})$-**Coalg**$(C,D)$ and $\theta' \in (\mathcal{A},\psi_{\mathcal{F}})$-**Coalg**$(C',D')$, we define $\theta\wedge\theta'$ to be the composite of (strict) right $\mathcal{A}$-module maps



where ι is induced by the intertwining map (0.2.1). It is straightforward to show that $\wedge$ endows $(\mathcal{A},\psi_{\mathcal{F}})$-**Coalg** with the structure of a monoidal category.

1. The general homotopy orbit model

Let Y be a topological space endowed with a left action of the circle S^1. The *homotopy orbit space* of Y, denoted Y_{hS^1}, is then $ES^1 \underset{S^1}{\times} Y$, where ES^1 is a contractible, free S^1-space.

In this section we explain how to construct a model for the cohomology algebra of Y_{hS^1} by twisting together $\mathrm{H}^*(BS^1)$ and CU^*Y. We begin by developing a framework in which to calculate the cohomology algebra of the quotient space of a general group action, using the operadic tools recalled in section 0.2. We then construct a particularly nice family of primitive elements in CU_*S^1, which we proceed to apply to building a highly structured resolution of CU_*ES^1. Using that resolution, we then define the desired model for Y_{hS^1} within the framework developed at the beginning of the section.

1.1 Homology of quotient spaces of group actions.

Let G be a connected topological group, and let E be a right G-space. We apply the operadic machinery recalled in section 0.2 to the definition of a very special type of resolution of CU_*E. We begin by defining appropriate compatibility between DCSH structure and multiplicative structure.

Definition. Suppose that H and K are chain Hopf algebras. A chain map $\theta : H \longrightarrow K$ is a *multiplicative DCSH map* if there is a map of right $\mathcal{A}$-modules

$$\widehat{\theta} : \mathcal{T}(H)\underset{\mathcal{A}}{\diamond}\mathcal{F} \longrightarrow \mathcal{T}(K)$$

such that $\widehat{\theta}(1)(x\otimes z_0) = \theta(x)$ for all $x\in H$ and such that

$$\begin{array}{ccc}
\mathcal{T}(H\wedge H)\underset{\mathcal{A}}{\diamond}\mathcal{F} & \xrightarrow{\mathcal{T}(\mu_H)\underset{\mathcal{A}}{\diamond}1} & \mathcal{T}(H)\underset{\mathcal{A}}{\diamond}\mathcal{F} \\
\downarrow{\scriptstyle \widehat{\theta}\wedge\widehat{\theta}} & & \downarrow{\scriptstyle \widehat{\theta}} \\
\mathcal{T}(K\wedge K) & \xrightarrow{\mathcal{T}(\mu_K)} & \mathcal{T}(K)
\end{array}$$

commutes, where μ_H and μ_K are the multiplication maps of H and K, which are maps of coalgebras.

A multiplicative DCSH map is necessarily a chain algebra map, as can be seen by looking at the diagram above in level 1.

Definition. Let $\theta : H \longrightarrow K$ be a multiplicative DCSH map. Let M and N be a right H-module coalgebra and a right K-module coalgebra, respectively, where ρ_M and ρ_N are the module structure maps. A chain map $\varphi : M \longrightarrow N$ is a *DCSH module map* with respect to θ if there is a map of right $\mathcal{A}$-modules

$$\widehat{\varphi} : \mathcal{T}(M) \underset{\mathcal{A}}{\diamond} \mathcal{F} \longrightarrow \mathcal{T}(N)$$

such that $\widehat{\varphi}(1)(y \otimes z_0) = \varphi(y)$ for all $y \in M$ and such that

$$\begin{array}{ccc} \mathcal{T}(M \otimes H) \underset{\mathcal{A}}{\diamond} \mathcal{F} & \xrightarrow{\mathcal{T}(\rho_M) \underset{\mathcal{A}}{\diamond} 1} & \mathcal{T}(M) \underset{\mathcal{A}}{\diamond} \mathcal{F} \\ \downarrow{\scriptstyle \widehat{\varphi} \wedge \widehat{\theta}} & & \downarrow{\scriptstyle \widehat{\varphi}} \\ \mathcal{T}(N \otimes K) & \xrightarrow{\mathcal{T}(\rho_N)} & \mathcal{T}(N) \end{array}$$

commutes. If M is a semifree extension of H (cf., section 0.1) and φ is a quasi-isomorphism, then φ is a *DCSH H-resolution* of N.

Examining the diagram above on level 1, we see that $\varphi\rho_M = \rho_N(\varphi \otimes \theta)$.

The definition of DCSH module maps of left module coalgebras can be deduced easily from the definition above.

Let H be a chain Hopf algebra. Suppose that M and M' are right and left H-module coalgebras, with structure maps ρ_M and $\lambda_{M'}$, respectively. Consider the following pushout of chain complexes.

$$\begin{array}{ccc} M \otimes H \otimes M' & \xrightarrow{1 \otimes \lambda_{M'}} & M \otimes M' \\ \downarrow{\scriptstyle \rho_M \otimes 1} & & \downarrow{\scriptstyle \bar{\rho}} \\ M \otimes M' & \xrightarrow{\bar{\lambda}} & M \underset{H}{\otimes} M' \end{array}$$

Since $\rho_M \otimes 1$ and $1 \otimes \lambda_{M'}$ are both maps of coalgebras, $M \underset{H}{\otimes} M'$ admits a coalgebra structure such that the quotient map

$$\pi = \bar{\lambda} = \bar{\rho} : M \otimes M' \to M \underset{H}{\otimes} M'$$

is a coalgebra map.

Theorem 1.1.1. *Let $\theta : H \longrightarrow K$ be a multiplicative DCSH map. Let M and M' be right and left H-module coalgebras, and let N and N' be right and left K-module coalgebras. Let $\varphi : M \longrightarrow N$ and $\varphi' : M' \longrightarrow N'$ be DCSH module maps with respect to θ. Then the induced chain map*

$$\varphi \underset{\theta}{\otimes} \varphi' : M \underset{H}{\otimes} M' \longrightarrow N \underset{K}{\otimes} N'$$

is a DCSH map. Furthermore, if in addition M' and N' are right L-comodule coalgebras, where L is a chain Hopf algebra, and φ' is a DCSH module map with respect to Id_L, then $\varphi \underset{\theta}{\otimes} \varphi'$ is a DCSH module map with respect to Id_L as well.

Proof. Colimits in $\mathbf{M}^{\Sigma}$ are calculated level-wise. Since $\mathcal{T}(A \wedge B)$ is naturally isomorphic to $\mathcal{T}(A) \otimes \mathcal{T}(B)$, it is easy to see that the diagrams


$$\begin{array}{ccc} \mathcal{T}(M \otimes H \otimes M') & \xrightarrow{\mathcal{T}(1\otimes\lambda_{M'})} & \mathcal{T}(M \otimes M') \\ \downarrow{\scriptstyle \mathcal{T}(\rho_M\otimes 1)} & & \downarrow{\scriptstyle \mathcal{T}(\pi)} \\ \mathcal{T}(M \otimes M') & \xrightarrow{\mathcal{T}(\pi)} & \mathcal{T}(M \underset{H}{\otimes} M') \end{array}$$

and

$$\begin{array}{ccc} \mathcal{T}(N \otimes K \otimes N') & \xrightarrow{\mathcal{T}(1\otimes\lambda_{N'})} & \mathcal{T}(N \otimes N') \\ \downarrow{\scriptstyle \mathcal{T}(\rho_N\otimes 1)} & & \downarrow{\scriptstyle \mathcal{T}(\pi)} \\ \mathcal{T}(N \otimes N') & \xrightarrow{\mathcal{T}(\pi)} & \mathcal{T}(N \underset{K}{\otimes} N') \end{array}$$

are pushouts in the category of right $\mathcal{A}$-modules, $\mathbf{Mod}_{\mathcal{A}}$. On the other hand, since the composition monoidal structure on $\mathbf{Mod}_{\mathcal{A}}$ is right closed, the endofunctor $- \underset{\mathcal{A}}{\diamond} \mathcal{F}$ is a left adjoint and therefore preserves pushouts.

Consider the following diagram.



The front and back sides are pushouts, and the left side and the top both commute since φ and φ' are DCSH module maps. Therefore, there exists a map of left $\mathcal{A}$-modules

$$\widehat{\varphi} \underset{\widehat{\theta}}{\wedge} \widehat{\varphi}' : \mathcal{T}(M \underset{H}{\otimes} M') \underset{\mathcal{A}}{\diamond} \mathcal{F} \longrightarrow \mathcal{T}(N \underset{K}{\otimes} N')$$

that makes the whole cube commute. Restricting to level 1, we verify easily that

$$\widehat{\varphi} \underset{\widehat{\theta}}{\wedge} \widehat{\varphi}'(1)(x \underset{H}{\otimes} x' \otimes z_0) = \widehat{\varphi}(x) \underset{K}{\otimes} \widehat{\varphi}'(x')$$

as desired.

From the diagram above, it is easy to see that if φ' is a DCSH module map with respect to Id_L, then $\varphi \underset{\theta}{\wedge} \varphi'$ is as well. $\square$

Let G be a connected topological group, let E be the total space of a principal G-bundle, where G acts on E on the right, and let Y be a left G-space. Let $r : E \times G \to E$ and $l : G \times Y \to Y$ be the actions. Let $p : E \times Y \to E \underset{G}{\times} Y$ denote the quotient map.

Recall that for any pair of spaces X and W, the Eilenberg-Zilber (or shuffle) equivalence $\mathrm{EZ} : CU_*X \otimes CU_*W \longrightarrow CU_*(X \times W)$ is a coalgebra map. Consequently, the induced maps

$$\begin{array}{ccccc} CU_*E \otimes CU_*G & \xrightarrow[\simeq]{\mathrm{EZ}} & CU_*(E \times G) & \xrightarrow{CU_*r} & CU_*X \\ & \searrow^{\rho} & & \nearrow & \end{array}$$

and

$$\begin{array}{ccccc} CU_*G \otimes CU_*Y & \xrightarrow[\simeq]{\mathrm{EZ}} & CU_*(G \times Y) & \xrightarrow{CU_*l} & CU_*Y \\ & \searrow^{\lambda} & & \nearrow & \end{array}$$

are coalgebra maps as well. In the following pushout of chain complexes, therefore,

$$\begin{array}{ccc} CU_*E \otimes CU_*G \otimes CU_*Y & \xrightarrow{1\otimes\lambda} & CU_*E \otimes CU_*Y \\ \downarrow{\scriptstyle \rho\otimes 1} & & \downarrow{\scriptstyle \pi} \\ CU_*E \otimes CU_*Y & \xrightarrow{\pi} & CU_*E \underset{CU_*G}{\otimes} CU_*Y, \end{array}$$

$CU_*E \underset{CU_*G}{\otimes} CU_*Y$ admits a coalgebra structure such that the quotient map π is a coalgebra map. Furthermore, the chain map $CU_*E \underset{CU_*G}{\otimes} CU_*Y \to CU_*(E \underset{G}{\times} Y)$ induced by the composite

$$CU_*E \otimes CU_*Y \xrightarrow{\mathrm{EZ}} CU_*(E \times Y) \xrightarrow{CU_*p} CU_*(E \underset{G}{\times} Y) \tag{1.1.1}$$

is actually a coalgebra map.

We now use the results above on DCSH resolutions to prove a more highly structured version of Moore's classic theorem [Mo].

Theorem 1.1.2. *Let $\theta : H \longrightarrow CU_*G$ be a multiplicative DCSH quasi-isomorphism. Let $\varphi : M \longrightarrow CU_*E$ be a DCSH H-resolution of CU_*E. Then there is a DCSH quasi-isomorphism*

$$M \otimes CU_*X \xrightarrow{\simeq} CU_*(E \underset{G}{\times} X).$$

In particular,

$$H^*\big((M \underset{H}{\otimes} CU_*X)^\sharp\big) \cong H^*(E \underset{G}{\times} X)$$

as graded algebras.

Proof. Recall that Moore proved in [Mo] that given any CU_*G-semifree resolution of CU_*E,

$$\psi : N \xrightarrow{\simeq} CU_*E$$

the composite

$$N \underset{CU_*G}{\otimes} CU_*X \xrightarrow{\psi\otimes Id} CU_*E \underset{CU_*G}{\otimes} CU_*X \to CU_*(E \underset{G}{\times} X)$$

is a quasi-isomorphism.

Observe now that according to Theorem 1.1.1, the induced map

$$\varphi \underset{\theta}{\wedge} Id_{CU_*G} : M \underset{H}{\otimes} CU_*G \to CU_*E \underset{CU_*G}{\otimes} CU_*G \cong CU_*E$$

is a DCSH module map with respect to Id_{CU_*G}.

For any chain algebra A, let $\mathcal{B}A\widetilde{\otimes}A$ denote the acyclic bar construction on A, which is an A-semifree resolution of the base ring R. Consider the composite

$$M \underset{\simeq}{\xrightarrow{\varphi}} CU_*E \underset{\simeq}{\xrightarrow{\text{incl.}}} CU_*E \otimes \mathcal{B}H\widetilde{\otimes}H.$$

Since it is a quasi-isomorphism of H-semifree modules, this composite induces a quasi-isomorphism

$$\alpha : M \underset{H}{\otimes} CU_*G \xrightarrow{\simeq} (CU_*E \otimes \mathcal{B}H\widetilde{\otimes}H) \underset{H}{\otimes} CU_*G,$$

by Proposition 2.3(i) in [FHT]. On the other hand, since $CU_*E \otimes \mathcal{B}H\widetilde{\otimes}H$ and $CU_*E \otimes \mathcal{B}CU_*G\widetilde{\otimes}CU_*G$ are both semifree, over H and CU_*G respectively, and θ is a quasi-isomorphism of chain algebras, Proposition 2.4 in [FHT] implies that the quasi-isomorphism

$$1 \otimes \mathcal{B}\theta \otimes \theta : CU_*E \otimes \mathcal{B}H\widetilde{\otimes}H \xrightarrow{\simeq} CU_*E \otimes \mathcal{B}CU_*G\widetilde{\otimes}CU_*G$$

induces a quasi-isomorphism

$$\beta : (CU_*E \otimes \mathcal{B}H\widetilde{\otimes}H) \underset{H}{\otimes} CU_*G \xrightarrow{\simeq} (CU_*E \otimes \mathcal{B}CU_*G\widetilde{\otimes}CU_*G) \underset{CU_*G}{\otimes} CU_*G.$$

Let

$$\pi : CU_*E \otimes \mathcal{B}CU_*G\widetilde{\otimes}CU_*G \xrightarrow{\simeq} CU_*E$$

denote the obvious projection.

It is obvious that $\varphi \underset{\theta}{\wedge} Id_{CU_*G} = \pi\beta\alpha$ and is therefore a quasi-isomorphism. Thus $\varphi \underset{\theta}{\wedge} Id_{CU_*G} : M \underset{H}{\otimes} CU_*G \to CU_*E$ is a DCSH CU_*G-resolution of CU_*E.

Consider next the composite

$$\begin{array}{ccc} M \underset{H}{\otimes} CU_*X & \xrightarrow{\cong} & (M \underset{H}{\otimes} CU_*G) \underset{CU_*G}{\otimes} CU_*X \\ & & \Big\downarrow {\scriptstyle (\varphi \underset{\theta}{\wedge} Id_{CU_*G}) \underset{Id_{CU_*G}}{\wedge} Id_{CU_*X}} \\ & & CU_*E \underset{CU_*G}{\otimes} CU_*X \\ & & \Big\downarrow \\ & & CU_*(E \underset{G}{\times} X), \end{array}$$

where the first map is an isomorphism of coalgebras and the last map is the strict coalgebra map (1.1.1). According to Theorem 1.1.1, the first vertical map is a DCSH map and its dual therefore induces an morphism of algebras in cohomology. Finally, by Moore's theorem, the composite of the two vertical maps is a quasi-isomorphism, so we can conclude. □

1.2 A special family of primitives.
In this section, as first step towards defining a DCSH resolution of CU_*ES^1, we identify an important family of primitive elements in CU_*S^1. We begin by defining a suspension-type degree +1 operation on CU_*S^1.

Definition. If $T : I^n \longrightarrow S^1$ is an n-cube, let $\sigma(T)$ be the $(n+1)$-cube defined by

$$\sigma(T)(t_0, ..., t_n) := \big(T(t_1, ..., t_n)\big)^{t_0},$$

where we are considering S^1 as the unit circle in the complex plane, i.e.,

$$S^1 = \{e^{i\theta} \mid 0 \leq \theta \leq 2\pi\}.$$

Remark. It is clear that $\sigma(T)$ is degenerate if T is degenerate. The operation σ can therefore be extended linearly to all of CU_*S^1.

As the next lemma states, σ is a contracting homotopy in degrees greater than one and is a $(Id, 0)$-coderivation.

Lemma 1.2.1. *Let $T \in CU_*S^1$.*

(1) *If $\deg T \geq 2$, then $d\sigma(T) = T - \sigma(dT)$ where d is the usual differential on CU_*S^1.*
(2) *$\overline{\Delta}(\sigma(T)) = \sigma(T_i) \otimes T^i$, where $\overline{\Delta}$ is the usual reduced coproduct on CU_*S^1 and $\Delta(T) = T_i \otimes T^i$ (using the Einstein summation convention).*

Simple calculations, applying the definitions of the cubical differential and the cubical coproduct, as given for example in [Mas] and [A], suffice to prove this lemma.

We now apply the σ operation to the recursive construction of an important family of elements in CU_*S^1.

Definition. Let $T_0 : I \longrightarrow S^1$ be the 1-cube defined by $T_0(t) = e^{i2\pi t}$. Given $T_k \in CU_{2k+1}S^1$ for all $k < n$, let T_n be the $(2n+1)$-cubical chain defined by

$$T_n := \sigma\left(\sum_{i=1}^{n} T_{i-1} \cdot T_{n-i}\right) \in CU_{2n+1}S^1.$$

Let $\mathfrak{T} := \{T_n \mid n \geq 0\}$.

Examples. It is easy to see that

$$T_1(t_0, t_1, t_2) = e^{i2\pi t_0(t_1+t_2)}$$

and that $T_2 = U + V$ where

$$U(t_0, ..., t_4) = e^{i2\pi t_0(t_1+(t_2+t_3)t_4)} \text{ and } V(t_0, ..., t_4) = e^{i2\pi t_0(t_1(t_2+t_3)+t_4)}.$$

Proposition 1.2.2. *The family $\mathfrak{T}$ satisfies the following properties.*

(1) *$dT_0 = 0$ and $0 \neq [T_0]$ in $H_1 S^1$.*
(2) *$dT_n = \sum_{i=1}^{n} T_{i-1} T_{n-i}$ for all $n > 0$.*
(3) *Every T_n is primitive in $CU_* S^1$.*

Proof. Points (1) and (2) are easy consequences of Lemma 1.2.1. It is well known that T_0 represents the unique nonzero homology generator of $H_* S^1$.

An easy inductive argument applying Lemma 1.2.1(2) proves point (3), since if T_k is primitive for all $k < n$, then the sum $\sum_{i=1}^{n} T_{i-1} \cdot T_{n-i}$ is also primitive, as it is symmetric and all factors are of odd degree. □

Let $\mathbb{T}$ denote the subalgebra of $CU_* S^1$ generated by the family $\mathfrak{T}$. Since all the T_n's are primitive, $\mathbb{T}$ is a sub Hopf algebra of $CU_* S^1$. Proposition 1.2.2(1) and (2) imply that $\mathbb{T}$ is closed under the differential.

It is very helpful to recognize $\mathbb{T}$ as the image of a certain homomorphism, as we next make explicit. Let Γ denote the divided powers algebra functor. If w is in even degree, then

$$\Gamma w = \bigoplus_{k \geq 0} \mathbb{Z} \cdot w(k),$$

where $\deg w(k) = k \cdot |w|$, $w(0) = 1$, $w(1) = w$ and $w(k)w(l) = \binom{k+l}{k} w(k+l)$. Furthermore, Γw is in fact a Hopf algebra, where the coproduct is specified by $\Delta(w) = w \otimes 1 + 1 \otimes w$. Recall that the integral homology $\mathrm{H}_* BS^1$ of the classifying space of the circle is isomorphic to Γv, where v is of degree 2.

Define a linear map $\zeta : s^{-1} \mathrm{H}_* BS^1 \to CU^* S^1$ by $\zeta(s^{-1} v(k)) = T_{k-1}$. A simple calculation, based on Proposition 1.2.2 (1) and (2), shows that ζ extends to a quasi-isomorphism of chain Hopf algebras

$$\widehat{\zeta} : \Omega \mathrm{H}_* BS^1 \xrightarrow{\simeq} CU_* S^1,$$

where $\Omega \mathrm{H}_* BS^1$ is primitively generated. It is clear that $\mathbb{T} = \mathrm{Im}\, \widehat{\zeta}$, so that, in particular, the inclusion

$$\mathbb{T} \hookrightarrow CU_* S^1$$

is a quasi-isomorphism.

1.3 A DCSH module resolution of $CU_* ES^1$.

Using the family $\mathfrak{T}$, we now construct a DCSH resolution of $CU_* ES^1$ as a $\Omega \mathrm{H}_* BS^1$-module.

Let $\mathrm{H}_* BS^1 \otimes_{t_\Omega} \Omega \mathrm{H}_* BS^1$ denote the acyclic cobar construction on $\mathrm{H}_* BS^1$ (cf., section 0.1). Explicitly, $\mathrm{H}_* BS^1 \otimes_{t_\Omega} \Omega \mathrm{H}_* BS^1 = (\Gamma v \otimes T s^{-1} \Gamma^+ v, D_\Omega)$, where

$$D_\Omega(v(n) \otimes w) = v(n) \otimes d_\Omega w - \sum_{i=0}^{n} v(i) \otimes s^{-1} v(n-i) \cdot w$$

for all n and for all $w \in T s^{-1} \Gamma^+ v$. It is easy to see that the usual tensor coproduct on $\Gamma v \otimes T s^{-1} \Gamma^+ v$ commutes with the differential, i.e., $\mathrm{H}_* BS^1 \otimes_{t_\Omega} \Omega \mathrm{H}_* BS^1$ is a chain coalgebra, with untwisted coproduct.

Let $j : S^1 \longrightarrow ES^1$ denote the inclusion of S^1 as the base of the construction of ES^1, which is an S^1-equivariant map. The composite

$$CU_* j \circ \widehat{\zeta} : \Omega \mathrm{H}_* BS^1 \longrightarrow CU_* ES^1$$

is map of $\Omega\,\mathrm{H}_*\,BS^1$-module coalgebras. Consider the following commutative diagram of right $\Omega\,\mathrm{H}_*\,BS^1$-module coalgebras.

$$\begin{array}{ccc} \Omega\,\mathrm{H}_*\,BS^1 & \xrightarrow{CU_*j\circ\widehat{\zeta}} & CU_*ES^1 \\ \downarrow{\scriptstyle \iota} & & \downarrow{\scriptstyle \simeq} \\ \mathrm{H}_*\,BS^1\otimes_{t_\Omega}\Omega\,\mathrm{H}_*\,BS^1 & \xrightarrow{\simeq} & \mathbb{Z} \end{array}$$

Since the inclusion ι is a coalgebra map, and therefore a DCSH map, and the composite $CU_*j\circ\widehat{\zeta}$ is a $\Omega\,\mathrm{H}_*\,BS^1$-module coalgebra map, and therefore a DCSH module map, Theorem 0.2.1 implies that they can be realized as morphisms of right $\mathcal{A}$-modules, as noted below.

$$\begin{array}{ccc} \mathcal{T}\big(\Omega\,\mathrm{H}_*\,BS^1\big)\underset{\mathcal{A}}{\diamond}\mathcal{F} & \xrightarrow{\ \mathfrak{j}\ } & \mathcal{T}\big(CU_*ES^1\big) \\ \downarrow{\scriptstyle \mathcal{T}(\iota)\underset{\mathcal{A}}{\diamond}1} & & \\ \mathcal{T}\big(\mathrm{H}_*\,BS^1\otimes_{t_\Omega}\Omega\,\mathrm{H}_*\,BS^1\big)\underset{\mathcal{A}}{\diamond}\mathcal{F} & & \end{array}$$

Since CU_*ES^1 is acyclic, $\mathrm{H}_*\,BS^1\otimes_{t_\Omega}\Omega\,\mathrm{H}_*\,BS^1$ is semifree, the underlying algebra of $\Omega\,\mathrm{H}_*\,BS^1$ is free and $\mathcal{F}$ is a semifree $\mathcal{A}$-bimodule, we can inductively construct an extension $\mathfrak{k}$ of $\mathfrak{j}$ so that

$$\begin{array}{ccc} \mathcal{T}\big(\Omega\,\mathrm{H}_*\,BS^1\big)\underset{\mathcal{A}}{\diamond}\mathcal{F} & \xrightarrow{\ \mathfrak{j}\ } & \mathcal{T}\big(CU_*ES^1\big) \\ \downarrow{\scriptstyle \mathcal{T}(\iota)\underset{\mathcal{A}}{\diamond}1} & \nearrow{\scriptstyle \mathfrak{k}} & \\ \mathcal{T}\big(\mathrm{H}_*\,BS^1\otimes_{t_\Omega}\Omega\,\mathrm{H}_*\,BS^1\big)\underset{\mathcal{A}}{\diamond}\mathcal{F} & & \end{array}$$

commutes. Furthermore, $\mathfrak{k}$ is necessarily a quasi-isomorphism, since both CU_*ES^1 and $\mathrm{H}_*\,BS^1\otimes_{t_\Omega}\Omega\,\mathrm{H}_*\,BS^1$ are acyclic. In other words, restricting to level 1, we obtain a DCSH $\Omega\,\mathrm{H}_*\,BS^1$-resolution of CU_*ES^1:

$$\mathrm{H}_*\,BS^1\otimes_{t_\Omega}\Omega\,\mathrm{H}_*\,BS^1\xrightarrow{\simeq}CU_*ES^1. \tag{1.3.1}$$

1.4 Modeling S^1-homotopy orbits.
Applying Theorem 1.1.2 to the DCSH resolution (1.3.1), we obtain a chain coalgebra model for S^1-homotopy orbits.

Theorem 1.4.1. *Let Y be any (left) S^1-space. Then there is a DCSH quasi-isomorphism*

$$\big(\mathrm{H}_*\,BS^1\otimes_{t_\Omega}\Omega\,\mathrm{H}_*\,BS^1\big)\underset{\Omega\,\mathrm{H}_*\,BS^1}{\otimes}CU_*Y\xrightarrow{\simeq}CU_*Y_{hS^1}.$$

In particular,

$$\mathrm{H}^*\left(\big((\mathrm{H}_*\,BS^1\otimes_{t_\Omega}\Omega\,\mathrm{H}_*\,BS^1)\underset{\Omega\,\mathrm{H}_*\,BS^1}{\otimes}CU_*Y\big)^\sharp\right)\cong\mathrm{H}^*(Y_{hS^1})$$

as graded algebras.

A careful examination of $(\mathrm{H}_* BS^1 \otimes_{t_\Omega} \Omega\,\mathrm{H}_* BS^1) \underset{\Omega\,\mathrm{H}_* BS^1}{\otimes} CU_*Y$ is certainly warranted. Let $g : S^1 \times Y \longrightarrow Y$ be the action map. Let κ denote the composite

$$CU_*S^1 \otimes CU_*Y \xrightarrow[\simeq]{\mathrm{EZ}} CU_*(S^1 \times Y) \xrightarrow{CU_*g} CU_*Y \ .$$

Recall that $\mathrm{H}_* BS^1 = \Gamma v$, where v is of degree two. Define an extension $(\Gamma v \otimes CU_*Y, D)$ of Γv by

$$D(v(n) \otimes U) := v(n) \otimes dU + \sum_{k=1}^{n-1} v(n-k-1) \otimes \kappa(T_k \otimes U).$$

By Proposition 1.2.2(2), $D^2 = 0$. Observe that $(\Gamma v \otimes CU_*Y, D)$ is naturally a chain coalgebra that is a cofree left Γv-comodule, since κ is a coalgebra map and each T_k is primitive.

It is easy to show that the following two maps are chain coalgebra isomorphisms, one inverse to the other.

$$(\mathrm{H}_* BS^1 \otimes_{t_\Omega} \Omega\,\mathrm{H}_* BS^1) \underset{\Omega\,\mathrm{H}_* BS^1}{\otimes} CU_*Y \longrightarrow (\Gamma v \otimes CU_*Y, D)$$

$$v(n) \otimes w \otimes U \longmapsto v(n) \otimes \kappa\big(\widehat{\zeta}(w) \otimes U\big)$$

$$(\Gamma v \otimes CU_*Y, D) \longrightarrow (\mathrm{H}_* BS^1 \otimes_{t_\Omega} \Omega\,\mathrm{H}_* BS^1) \underset{\Omega\,\mathrm{H}_* BS^1}{\otimes} CU_*Y$$

$$v(n) \otimes U \longmapsto v(n) \otimes 1 \otimes U$$

Composing the second isomorphism with the DCSH quasi-isomorphism of Theorem 1.4.1, we obtain a DCSH quasi-isomorphism

$$(\Gamma v \otimes CU_*Y, D) \longrightarrow CU_*Y_{hS^1} \tag{1.4.1}$$

that fits into the following commutative diagram.

$$\begin{array}{ccccc} CU_*Y & \xrightarrow{\text{incl.}} & (\Gamma v \otimes CU_*Y, D) & \xrightarrow{\text{proj.}} & (\Gamma v, 0) \\ \downarrow{=} & & \downarrow{\simeq} & & \downarrow{\simeq} \\ CU_*Y & \longrightarrow & CU_*Y_{hS^1} & \longrightarrow & CU_*BS^1 \end{array}$$

In this article we are interested in cohomology calculations and so must dualize this model. Let $(\Lambda v \otimes CU^*Y, D^\sharp)$ denote the R-dual of $(\Gamma v \otimes CU_*X, D)$. In particular $v(v) = 1$. Since it is the dual of a cofree comodule, $(\Lambda v \otimes CU^*X, D^\sharp)$ is a free, right Λv-module. The dual of (1.4.1)

$$CU^*(Y_{hS^1}) \longrightarrow (\Lambda v \otimes CU^*Y, D^\sharp) \tag{1.4.2}$$

is a quasi-isomorphism inducing an algebra map in cohomology, which fits into a commutative diagram

$$\begin{array}{ccccc} (\Lambda v, 0) & \xrightarrow{\text{incl.}} & (\Lambda v \otimes CU^*Y, D^\sharp) & \xrightarrow{\text{proj.}} & CU^*Y \\ {\scriptstyle =}\uparrow & & \uparrow{\scriptstyle \simeq} & & \uparrow{\scriptstyle \simeq} \\ CU^*BS^1 & \longrightarrow & CU^*Y_{hS^1} & \longrightarrow & CU^*Y \end{array} \tag{1.4.3}$$

This is the *cubical S^1-homotopy orbit model*, which we denote $\mathrm{HOS}^*(Y)$.

A simple dualization calculation enables us to describe $D^\sharp$ completely. For each $n \geq 0$, define $\omega_n : CU^*Y \longrightarrow CU^{*-(2n+1)}Y$ to be the R-dual of $\kappa(T_n \otimes -)$.

Lemma 1.4.2. *If $f \in CU^mY$, then*

$$D^\sharp(v^n \otimes f) = v^n \otimes d^\sharp f + \sum_{k=0}^{\lceil \frac{m-1}{2} \rceil} v^{n+k+1} \otimes \omega_k(f)$$

where $d^\sharp$ denotes the differential of CU^*Y.

As a consequence of this description of $D^\sharp$, we obtain the following useful properties of the operators ω_k.

Corollary 1.4.3. *The operators ω_n satisfy the following properties.*

1. *For all $n \geq 1$, $[d^\sharp, \omega_n] = -\sum_{k=0}^{n-1} \omega_k \circ \omega_{n-k-1}$, while $d^\sharp\omega_0 = -\omega_0 d^\sharp$.*
2. *Each ω_n is a derivation, i.e., $\omega_n(f \cdot g) = \omega_n(f) \cdot g + (-1)^f f \cdot \omega_n(g)$.*

Proof. The proof of (1) proceeds by expansion of the equation $0 = (D^\sharp)^2(1 \otimes f)$. To prove (2), expand the equation

$$D^\sharp(1 \otimes f \cdot g) = D^\sharp(1 \otimes f) \cdot (1 \otimes g) + (-1)^f (1 \otimes f) \cdot D^\sharp(1 \otimes g).$$

The differential $D^\sharp$ is a derivation, since it is the dual of the differential of a chain coalgebra. □

Remark. This corollary implies that ω_0 induces a derivation of degree -1

$$\varpi : \mathrm{H}^* Y \longrightarrow \mathrm{H}^{*-1} Y$$

such that $\varpi^2 = 0$. Let $\mathcal{C}_2$ denote the topological "little squares" operad, the homology of which is equivalent to the Gerstenhaber operad $\mathcal{G}$ governing Gerstenhaber algebras. It is well known that $\mathcal{C}_2(2)$ is homotopy equivalent to S^1, so that the generator of $\mathrm{H}_1 S^1$ corresponds to the Gerstenhaber bracket operation [K]. The derivation ϖ must therefore be closely related to the Gerstenhaber bracket, since a representative of the generator of $\mathrm{H}_1 S^1$ gives rise to it. It is in fact the Δ-operation of the Batalin-Vilkovisky structure on $\mathrm{H}^* X$ [G].

2. S^1-homotopy orbits of the free loop space

In section 1 we constructed a cochain algebra model $\mathrm{HOS}^*(Y)$ for homotopy orbits on an S^1-space Y, by twisting together $\mathrm{H}^* BS^1$ and CU^*Y. The model we obtained is quite large, however, since the cubical cochains on Y are not even of finite type, if Y is not a finite or discrete space. In this section we show how to reduce the model of section 1 even further when Y is a free loop space, obtaining a model with which it is possible to make explicit calculations. To do so, we replace the cubical cochains on Y by an equivalent and much smaller cochain complex, derived from the free loop space model of [DH] and [H].

We begin this section by recalling the construction of [DH] and [H], which provides us with an associative cochain algebra that is weakly equivalent to the normalized cochains on a certain simplicial model of the free loop space. We then explain how to relate this model to cubical cochains. Finally, we "linearize" the free loop space model, obtaining a quasi-isomorphism of cochain complexes $\Upsilon : fls^*(X) \longrightarrow CU^*(\mathcal{L}X)$. We include only those proofs that have not already appeared in [H].

Restricting to an interesting class of spaces that includes all double suspensions, we "twist" together $fls^*(X)$ and H^*BS^1, obtaining a cochain complex $hos^*(X)$ such that $\mathrm{H}^*(hos^*(X)) \cong \mathrm{H}^*(\mathcal{L}X_{hS^1})$ as graded $\mathbb{Z}$-modules. We then show that for any space X in the class we consider and over any field $\Bbbk$, the free loop model map Υ lifts to a quasi-isomorphism $\widetilde{\Upsilon} : hos^*(X) \longrightarrow CU^*(\mathcal{L}X_{hS^1})$.

To conclude this section we apply the $hos^*(-)$ construction to computing the cohomology of the S^1-homotopy orbits of $\mathcal{L}S^n$, as an illustration of relative ease of computation with such a small model. The Poincaré series of this equivariant cohomology was already known [BO].

The crucial building block in all of our constructions comes from [HPST]. In [HPST] the authors proved that for every 1-reduced simplicial set K, there is a natural, coassociative coproduct ψ_K on ΩC_*K with respect to which Szczarba's natural equivalence of chain algebras $\theta_K : \Omega C_*K \longrightarrow C_*GK$ is a DCSH map, i.e., there is a morphism of chain algebras $\Omega^2 C_*K \longrightarrow \Omega C_*GK$ of which $s^{-1}\theta_K s$ is the linear piece (see section 0.2). They established moreover that ψ_K is identical to the coproduct defined purely combinatorially by Baues in [B]. As a consequence, we know that

$$\operatorname{Im}\psi_K \subseteq Ts^{-1}C_+K \otimes s^{-1}C_+K. \tag{2.1}$$

The R-dual of ψ_K, denoted $\psi_K^\sharp$, is a natural multiplication on $\mathcal{B}C^*K$, which is the R-dual of ΩC_*K, since K is of finite type. Because of (2.1),

$$\perp sC^+K \otimes \perp^{>1} sC^+K \subseteq \ker \pi\psi_K^\sharp,$$

where $\pi : \perp sC^+K \longrightarrow sC^+K$ is the projection, i.e., the linear term of $\psi_K^\sharp(w \otimes sx_1|...|sx_n)$ is necessarily 0 for all $n > 1$. In particular, if $\psi_K^\sharp$ is commutative, then the linear term of $\psi_K^\sharp(sy_1|\cdots|sy_m \otimes sx_1|...|sx_n)$ is necessarily 0 if $m > 1$ or $n > 1$.

Throughout this section, we work only with 1-reduced simplicial sets K such that $\psi_K^\sharp$ is commutative. As follows from the chain-level Bott-Samelson theorem in [HPS2], if the simplicial set K is such that C_*K is cocommutative, then $\psi_{EK}^\sharp$ is commutative, where EK denotes the simplicial suspension of K [Ma]. In particular, if all elements of C_*K are primitive, e.g., if $C_nK = 0$ for all $n < r$ and for all $n \geq 2r$ for some r or if K is itself a suspension, then $\psi_{EK}^\sharp$ is commutative.

2.1 The simplicial and cubical free loop space models.

Let K be a 1-reduced simplicial set of finite type. Recall that the unit η of the $(|\cdot|, \mathcal{S}_\bullet)$-adjunction induces an injective quasi-isomorphism of chain coalgebras

$$C_*\eta_K : C_*K \xrightarrow{\simeq} S_*|K|$$

that dualizes to a natural, surjective quasi-isomorphism of cochain algebras

$$C^*\eta_K : S^*|K| \xrightarrow{\simeq} C^*K\ ,$$

where $S_* = C_* \circ \mathcal{S}_\bullet$ and $S^* = C^* \circ \mathcal{S}_\bullet$.

The *canonical free loop construction on* K, as defined in [H], is the twisted cartesian product $\mathcal{L}K = GK \underset{\bar\tau}{\times} K$, with structure group $GK \times GK$, where $GK \times GK$ acts on GK by $(v, w) \cdot u := vuw^{-1}$ and

$$\bar\tau : K \longrightarrow GK \times GK : x \longmapsto (\bar x, \bar x)\,.$$

It is shown in the proof of Proposition 1.1.1 in [H] that there is a natural trivial cofibration of simplicial sets $\hat\eta : \mathcal{L}K \xrightarrow{\simeq} \mathcal{S}_\bullet\mathcal{L}|K|$ that fits into a commutative diagram of morphisms of simplicial sets

$$\begin{array}{ccc} \mathcal{L}K & \xrightarrow{q} & K \\ {\scriptstyle\hat\eta}\downarrow{\scriptstyle\simeq} & & {\scriptstyle\eta}\downarrow{\scriptstyle\simeq} \\ \mathcal{S}_\bullet\mathcal{L}|K| & \xrightarrow{e} & |K|, \end{array}$$

where q is the projection and e is the free loop fibration map.

A more general version of the theorem below is proved in [Bl], for arbitrary 1-reduced K of finite type. Since in any case our construction of the homotopy orbit model requires that $\psi^\sharp_K$ be commutative, we state only the special case here.

Henceforth, we let $a \star b$ denote the product $\psi^\sharp_K(a \otimes b)$ for all $a, b \in \mathcal{B}C^*K$.

Theorem 2.1.1 [DH], [H]**.** *Let K be a 1-reduced simplicial set of finite type such that the natural product $\psi^\sharp_K$ on $\mathcal{B}C^*K$ is commutative. There is a twisted algebra extension*

$$\iota : \Omega\mathcal{B}C^*K \longrightarrow \Omega\mathcal{B}C^*K \odot \mathcal{B}C^*K$$

and a quasi-algebra quasi-isomorphism

$$\delta : \Omega\mathcal{B}C^*K \odot \mathcal{B}C^*K \xrightarrow{\simeq} C^*(\mathcal{L}K)$$

such that

$$\begin{array}{ccc} \Omega\mathcal{B}C^*K & \xrightarrow{\iota} & \Omega\mathcal{B}C^*K \odot \mathcal{B}C^*K \\ {\scriptstyle\varepsilon_K}\downarrow{\scriptstyle\simeq} & & {\scriptstyle\delta}\downarrow{\scriptstyle\simeq} \\ C^*K & \xrightarrow{C^*q} & C^*\mathcal{L}K \end{array}$$

commutes, where $\varepsilon_K : \Omega\mathcal{B}C^*K \to C^*K$ denotes the counit of the cobar-bar adjunction. Moreover, the algebra extension satisfies the following conditions.

(1) *The left action of $\Omega\mathcal{B}C^*K$ is specified recursively by*

$$\begin{aligned}\big(s^{-1}a\otimes 1\big)\cdot(1\otimes sx_1|\cdots|sx_n)&\\ =&s^{-1}a\otimes sx_1|\cdots|sx_n\\ &-\sum_{j=1}^{n-1}\Big[s^{-1}\big(a\star(sx_1|\cdots|sx_j)\big)\cdot(1\otimes sx_{j+1}|\cdots|sx_n)\\ &\qquad+(-1)^{(\theta_j)(\theta_n-\theta_j)}s^{-1}\big(a\star(sx_{j+1}|\cdots|sx_n)\big)\otimes sx_1|\cdots|sx_j\Big]\end{aligned}$$

where $\theta_j = j+\sum_{i\le j}|x_i|$.

(2) *The restriction of the differential $\overline{D}$ of the twisted extension to $1\otimes\mathcal{B}C^*K$ is specified recursively by*

$$\begin{aligned}\overline{D}(1\otimes sx_1|\cdots|sx_n) =&1\otimes d_{\mathcal{B}}(sx_1|\cdots|sx_n)\\ &+\sum_{j=1}^{n-1}\big[(s^{-1}(sx_1|\cdots|sx_j)\otimes 1)\cdot(1\otimes sx_{j+1}|\cdots|sx_n)\\ &\qquad-(-1)^{\theta_j(\theta_n-\theta j)}s^{-1}(sx_{j+1}|\cdots|sx_n)\otimes sx_1|\cdots|sx_j\big]\end{aligned}$$

with θ_j as above.

(3) *Let $a = sx_1|\cdots|sx_m, b = sy_1|\cdots|sy_n \in \mathcal{B}C^*K$. Then*

$$(1\otimes a)\cdot(1\otimes b) := 1\otimes a\star b.$$

The definitions of twisted algebra extensions and of quasi-algebra maps are recalled in section 0.1.

We now adapt the free loop model above to the case of topological spaces, in order to facilitate construction of a model of the S^1-homotopy orbits on the free loop space. More precisely, if X has the homotopy type of $|K|$, we show that we can link the model above directly to $CU^*(\mathcal{L}X)$.

Let X be a 1-connected space with the homotopy type of a finite-type CW-complex, and let K be a finite-type, 1-reduced simplicial set such that $|K|\simeq X$.

Since the underlying graded abelian group of C^*K is free, the surjective quasi-isomorphism of cochain algebras $C^*\eta : S^*|K| \to C^*K$ admits a cochain section $\mathfrak{s} : C^*K \to S^*|K|$ inducing an isomorphism of algebras in cohomology. Having fixed $\mathfrak{s}$, consider the following commutative diagram.

$$\begin{array}{ccccc} C^*K & \xrightarrow{\mathfrak{s}} & S^*|K| & \xrightarrow{S^*e} & S^*\mathcal{L}|K| \\ {\scriptstyle C^*q}\downarrow & & & & \downarrow{\scriptstyle C^*\hat{\eta}}\ \simeq \\ C^*\mathcal{L}K & & \xrightarrow{\quad=\quad} & & C^*\mathcal{L}K \end{array}$$

Since C^*q is injective and $C^*\hat{\eta}$ is a surjective quasi-isomorphism, $C^*\hat{\eta}$ admits a section $\hat{\mathfrak{s}}$ such that

$$\begin{array}{ccc} C^*K & \xrightarrow{C^*q} & C^*\mathcal{L}K \\ {\scriptstyle \mathfrak{s}}\downarrow{\scriptstyle \simeq} & & {\scriptstyle \hat{\mathfrak{s}}}\downarrow{\scriptstyle \simeq} \\ S^*|K| & \xrightarrow{S^*e} & S^*\mathcal{L}|K| \end{array}$$

commutes.

Recall now that for all spaces Y, subdivision of cubes defines a natural quasi-isomorphism of chain coalgebras

$$\xi : CU_*Y \xrightarrow{\simeq} S_*Y,$$

which, upon dualization, gives rise to a natural quasi-isomorphism of cochain algebras

$$\xi^\sharp : S^*Y \xrightarrow{\simeq} CU^*Y.$$

There is therefore a commuting diagram

$$\begin{array}{ccc} \Omega\mathcal{B}C^*K & \xrightarrow{\iota} & \Omega\mathcal{B}C^*K \odot \mathcal{B}C^*K \\ {\scriptstyle \xi^\sharp \mathfrak{s}\varepsilon_K}\downarrow{\scriptstyle \simeq} & & {\scriptstyle \xi^\sharp \hat{\mathfrak{s}}\delta}\downarrow{\scriptstyle \simeq} \\ CU^*X & \xrightarrow{CU^*e} & CU^*\mathcal{L}X, \end{array} \tag{2.1.1}$$

where the vertical arrows are quasi-algebra quasi-isomorphisms.

2.2 Linearization of the free loop model.

In this section we simplify the free loop model, making it as small as possible, to facilitate homotopy orbit space computations in section 2.4. Recall that we are considering only the case in which the natural product $\psi_K^\sharp$ on $\mathcal{B}C^*K$ is strictly commutative. The construction presented here is an improved version of that in [H].

Consider the surjection

$$\varepsilon_K \otimes Id : \Omega\mathcal{B}C^*K \otimes \mathcal{B}C^*K \longrightarrow C^*K \otimes \mathcal{B}C^*K.$$

Extend the differential on C^*K to a differential $\check{D}$ on $C^*K \otimes \mathcal{B}C^*K$, which is a free right C^*K-module, by

$$\check{D}(1 \otimes c) := (\varepsilon_K \otimes Id)\overline{D}(1 \otimes c),$$

extended as a right module derivation. Consequently, $\check{D}(\varepsilon_K \otimes Id) = (\varepsilon_K \otimes Id)\overline{D}$, which implies in turn that $\check{D}^2 = 0$.

Let $\pi : \mathcal{B}C^*K \longrightarrow C^*K$ be the projection onto (desuspended) linear terms, i.e.,

$$\pi(sx_1|\cdots|sx_n) := \begin{cases} x_1 : n = 1 \\ 0 : n > 1. \end{cases}$$

Recall from the discussion in the introduction to section 2 that since $\psi_K^\sharp$ is strictly commutative,

$$\pi\psi_K^\sharp(sy_1|\cdots|sy_m\otimes sx_1|\cdots|sx_n)=0 \quad \text{if } m>1 \text{ or } n>1. \tag{2.2.1}$$

Thus, from the definition of $\Omega\mathcal{B}C^*K\odot\mathcal{B}C^*K$ in Theorem 2.1.1, we obtain the following explicit formula for $\check{D}$, when $c=sx_1|\cdots|sx_n$.

$$\begin{aligned}\check{D}(1\otimes c)=&1\otimes d_{\mathcal{B}}c\\ &+\sum_{i\le n}\pi\big(sx_1\star\cdots\star sx_{i-1}\big)\otimes sx_i|\cdots|sx_n\\ &+\sum_{i<n}\pm\pi\big(sx_1\star\cdots\star sx_{i-1}\star sx_n\big)\otimes sx_i|\cdots|sx_{n-1},\end{aligned} \tag{2.2.2}$$

where the signs in the third line arise from applying the Koszul rule. Notice that it is entirely possible that $\check{D}(1\otimes c)\neq 1\otimes d_{\mathcal{B}}c$. On the other hand, if $\psi_K^\sharp$ is the shuffle product, then $\check{D}(1\otimes c)=1\otimes d_{\mathcal{B}}c$ for all $c\in\perp sC^+K$.

Define now a left C^*K-action on $C^*K\otimes\mathcal{B}C^*K$ by letting

$$(x\otimes 1)\cdot(1\otimes c):=(\varepsilon_K\otimes Id)\big((s^{-1}(sx)\otimes 1)\cdot(1\otimes c)\big)$$

for all $x\in C^*K$ and $c\in\mathcal{B}C^*K$, then extending to a morphism of free right C^*K-modules. If

$$\overline{D}(1\otimes c)-1\otimes d_{\mathcal{B}}c=\sum_i s^{-1}(a_i)\otimes b^i,$$

then

$$\begin{aligned}&\check{D}\big((x\otimes 1)\cdot(1\otimes c)\big)\\ &=(\varepsilon_K\otimes Id)\big((s^{-1}(sdx)\otimes 1)\cdot(1\otimes c)+(-1)^x(s^{-1}(sx)\otimes 1)\cdot\overline{D}(1\otimes c)\big)\\ &=(dx\otimes 1)\cdot(1\otimes c)\\ &\quad+(-1)^x(\varepsilon_K\otimes Id)\bigg((s^{-1}(sx)\otimes 1)\cdot\big(1\otimes d_{\mathcal{B}}c\\ &\qquad\qquad+\sum_i(-1)^{(a_i+1)b^i}(1\otimes b^i)(s^{-1}(sa_i)\otimes 1)\big)\bigg)\\ &=(dx\otimes 1)\cdot(1\otimes c)\\ &\quad+(-1)^x\bigg((x\otimes 1)\cdot\big(1\otimes d_{\mathcal{B}}c+\sum_i(-1)^{(a_i+1)b^i}(1\otimes b^i)(a_i\otimes 1)\big)\bigg)\\ &=(dx\otimes 1)\cdot(1\otimes c)+(-1)^x(x\otimes 1)\cdot\check{D}(1\otimes c),\end{aligned}$$

i.e., the left C^*K action commutes with the differential.

Using the commutativity of $\psi_K^\sharp$, for any $y\in C^*K$ and $c=sx_1|\cdots|sx_n\in\mathcal{B}C^*K$, we can write

$$\begin{aligned}(y\otimes 1)\cdot(1\otimes c)=&y\otimes c\\ &-\sum_i\pi\big(sy\star sx_1\star\cdots\cdots sx_{i-1}\big)\otimes sx_i|\cdots|sx_n\\ &-\sum_i\pm\pi\big(sy\star sx_1\star\cdots\cdots sx_{i-1}\star sx_n\big)\otimes sx_i|\cdots|sx_{n-1},\end{aligned} \tag{2.2.3}$$

where the signs in the third line arise from applying the Koszul rule. Note that the formulas above imply that when $\psi_K^\sharp$ is the shuffle product, the left action of C^*K is untwisted, i.e., $(x\otimes 1)(1\otimes c)=x\otimes c$.

It is easy to see that the left C^*K-action is defined precisely so that $\varepsilon_K\otimes Id$ is a map of left $\Omega\mathcal{B}C^*K$-modules. Since the right actions of $\Omega\mathcal{B}C^*K$ on $\Omega\mathcal{B}C^*K\odot\mathcal{B}C^*K$ and of C^*K on $C^*K\odot\mathcal{B}C^*K$ are both free, $\varepsilon_K\otimes Id$ is a map of right $\Omega\mathcal{B}C^*k$-modules as well. Consequently, for all $c,c'\in\mathcal{B}C^*K$,

$$\begin{aligned}\check{D}(1\otimes c\star c') =&(\varepsilon_K\otimes Id)\overline{D}(1\otimes c\star c')\\ =&(\varepsilon_K\otimes Id)\big[\overline{D}(1\otimes c)\cdot(1\star c')+(-1)^c(1\otimes c)\cdot\overline{D}(1\star c')\\ =&\check{D}(1\otimes c)\cdot(1\otimes c')+(-1)^c(1\otimes c)\cdot\check{D}(1\otimes c'),\end{aligned}$$

i.e., we can set

$$(1\otimes c)\cdot(1\otimes c')=1\otimes c\star c', \tag{2.2.4}$$

obtaining a differential algebra structure on $C^*K\otimes\mathcal{B}C^*K$, with respect to which $\varepsilon_K\otimes Id$ is an algebra map.

The following theorem summarizes the observations above.

Theorem 2.2.1. *Let $C^*K\odot\mathcal{B}C^*K$ denote $C^*K\otimes\mathcal{B}C^*K$ endowed with the differential $\check{D}$ of (2.2.2) and the multiplicative structure of (2.2.3) and (2.2.4). There is a twisted algebra extension*

$$C^*K\longrightarrow C^*K\odot\mathcal{B}C^*K\longrightarrow\mathcal{B}C^*K$$

such that $\varepsilon_K\otimes Id:\Omega\mathcal{B}C^*K\odot\mathcal{B}C^*K\longrightarrow C^*K\odot\mathcal{B}C^*K$ is a quasi-isomorphism of chain algebras.

For the constructions in section 2.3, we need a quasi-isomorphism

$$\Upsilon: C^*K\odot\mathcal{B}C^*K\xrightarrow{\simeq}CU^*\mathcal{L}X\,,$$

which we obtain as follows. Recall that ε_K has a differential, though not multiplicative, section

$$\sigma_K: C^*K\longrightarrow\Omega\mathcal{B}C^*K: x\longmapsto s^{-1}(sx).$$

Consider the following commutative diagram of cochain complexes and maps.

$$\begin{array}{ccccc} C^*K\odot\perp^{\leq 2}sC^+K & \xrightarrow{\sigma_K\otimes Id} & \Omega\mathcal{B}C^*K\odot\perp^{\leq 2}sC^+K & \xrightarrow{\text{incl.}} & \Omega\mathcal{B}C^*K\odot\mathcal{B}C^*K\\ \downarrow{\scriptstyle\text{incl.}} & & & & \simeq\downarrow{\scriptstyle\varepsilon_K\otimes Id}\\ C^*K\odot\mathcal{B}C^*K & & \xrightarrow{\qquad\qquad Id\qquad\qquad} & & C^*K\widetilde{\otimes}\mathcal{B}C^*K\end{array}$$

Since the inclusion map on the left is a free extension of cochain complexes and $\varepsilon_K\otimes Id$ is a surjective quasi-isomorphism, we can extend $\sigma_K\otimes Id$ to a cochain map

$$\hat{\sigma}: C^*K\odot\mathcal{B}C^*K\longrightarrow\Omega\mathcal{B}C^*K\odot\mathcal{B}C^*K$$

such that $(\varepsilon_K \otimes Id)\hat{\sigma} = Id$, i.e., $\hat{\sigma}$ is a section of $\varepsilon_K \otimes Id$. In particular, $\hat{\sigma}$ is a quasi-isomorphism and for all $x \otimes c \in C^*K \otimes \mathcal{B}C^*K$

$$s^{-1}(sx) \otimes c - \hat{\sigma}(x \otimes c) \in \ker \varepsilon_K \otimes \mathcal{B}C^*K.$$

Furthermore, since $\varepsilon_K \otimes Id$ is a chain algebra map inducing an isomorphism in cohomology, $\mathrm{H}^*\hat{\sigma} = \mathrm{H}^*(\varepsilon_K \otimes Id)^{-1}$ is an isomorphism of graded algebras. In particular, $\hat{\sigma}$ is a quasi-algebra map.

We now define Υ to be the composition below, where the map $\xi^\sharp \hat{\mathfrak{s}} \delta$ comes from diagram (2.1.1).

$$\begin{array}{ccccc} C^*K \odot \mathcal{B}C^*K & \xrightarrow{\hat{\sigma}} & \Omega\mathcal{B}C^*K \odot \mathcal{B}C^*K & \xrightarrow{\xi^\sharp \hat{\mathfrak{s}} \delta} & CU^*\mathcal{L}X \\ & \searrow & & \nearrow_{\Upsilon} & \end{array}$$

We have therefore constructed a commutative diagram

$$\begin{array}{ccc} C^*K & \xrightarrow{\iota} & C^*K \odot \mathcal{B}C^*K \\ {\scriptstyle\gamma}\downarrow{\scriptstyle\simeq} & & {\scriptstyle\Upsilon}\downarrow{\scriptstyle\simeq} \\ CU^*X & \xrightarrow{CU^*e} & CU^*\mathcal{L}X, \end{array} \tag{2.2.4}$$

where $\gamma = \xi^\sharp \mathfrak{s} \varepsilon_K \sigma_K$ (cf., diagram (2.1.1)).

Definition. Let X be a 1-connected space with the homotopy type of a finite-type CW-complex, and let K be a finite-type, 1-reduced simplicial set such that $|K| \simeq X$. Suppose that the natural product $\psi_K^\sharp$ on $\mathcal{B}C^*K$ is commutative. The twisted algebra extension

$$fls^*(X) := C^*K \odot \mathcal{B}C^*K$$

of Theorem 2.2.1, together with the quasi-algebra quasi-isomorphism

$$\Upsilon : fls^*(X) \xrightarrow{\simeq} CU^*\mathcal{L}X$$

of diagram (2.2.4), is a *thin free loop model* for X.

Since the Hochschild homology of C^*K is isomorphic to the cohomology of $CU^*\mathcal{L}X$, as shown in [J], it is interesting to compare the Hochschild complex of C^*K with the thin free loop model. As stated in the next proposition, the thin free loop model and the Hochschild complex on C^*K are isomorphic, at least in the special case where the multiplication on C^*K is trivial and $\psi_K^\sharp$ is commutative. This observation first arose in discussions several years ago with N. Dupont.

We have two reasons for choosing to work with the thin free loop model instead of the Hochschild complex. First, the thin free loop model is linked via a specific quasi-algebra quasi-isomorphism with $CU^*\mathcal{L}X$. To model constructions derived from the free loop space, such as its homotopy orbits, it is essential to have such a direct link with the cochains on $\mathcal{L}X$. Second, the cohomology of the thin free loop model is isomorphic as a graded algebra to the cohomology of $\mathcal{L}X$, whereas the isomorphism with Hochschild homology of C^*K is only an isomorphism of graded modules.

Proposition 2.2.2. *Let K be a 1-reduced, finite-type simplicial set such that the multiplication on C^*K is trivial and $\psi_K^\sharp$ is commutative. Let $\mathfrak{H}(C^*K)$ denote the Hochschild complex of C^*K. Then there is an isomorphism of cochain complexes*

$$\mathfrak{h} : C^*K \odot \mathcal{B}C^*K \xrightarrow{\cong} \mathfrak{H}(C^*K).$$

Proof. Let (A, d) be a cochain algebra. Recall that the Hochschild complex on (A, d) is

$$\mathfrak{H}(A, d) := (A \otimes \perp sA^+, D_{\mathfrak{H}})$$

where

$$\begin{aligned} D_{\mathfrak{H}}(a \otimes sb_1|\cdots|sb_n) =& da \otimes sb_1|\cdots|sb_n + (-1)^a a \otimes d_{\mathcal{B}}(sb_1|\cdots|sb_n) \\ & - ab_1 \otimes sb_2|\cdots|sb_n - (-1)^\epsilon b_n a \otimes sb_1|\cdots|sb_{n-1}, \end{aligned}$$

$d_{\mathcal{B}}$ is the usual bar construction differential and ϵ is the sign exponent dictated by the Koszul rule. In particular, if the multiplication in A is trivial, then $\mathfrak{H}(A, d) = (A, d) \otimes \mathcal{B}(A, d)$.

Define an endomorphism $\mathfrak{h}$ of the graded module $C^*K \otimes \perp sC^+K$ by

$$\mathfrak{h}(y \otimes 1) = y \otimes 1,$$

$$\mathfrak{h}(y \otimes sx_1|\cdots|sx_n) = y \otimes sx_1|\cdots|sx_n + x_n \otimes s\pi(sy \star sx_1)|\cdots|sx_{n-1},$$

for all $y \in C^+K$, and

$$\begin{aligned} \mathfrak{h}(1 \otimes sx_1|\cdots|sx_n) =& 1 \otimes sx_1|\cdots|sx_n \\ & + \sum_{j>1} 1 \otimes s\pi(sx_1 \star \cdots \star sx_j)|sx_{j+1}|\cdots|sx_n \\ & - \sum_{j\geq 1} \pm 1 \otimes s\pi(sx_n \star sx_1 \star \cdots \star sx_j)|sx_{j+1}|\cdots|sx_{n-1}, \end{aligned}$$

where the signs in the third line arise upon applying the Koszul rule. It is clear that $\mathfrak{h}$ is injective and therefore an isomorphism, since $C^*K \otimes \perp sC^+K$ is a free graded module of finite type.

Observe that (2.2.1) implies that

$$\pi\big(s\pi(sw_1 \star \cdots \star sw_m) \star sz\big) = \pi(sw_1 \star \cdots \star sw_m \star sz)$$

for all $w_1, ..., w_m, z \in C^*K$. Straightforward calculations based on this observation, as well as on the fact that C^*K is a trivial algebra, show that $D_{\mathfrak{H}}\mathfrak{h} = \mathfrak{h}\check{D}$. $\square$

Note that the isomorphism $\mathfrak{h}$ endows $\mathfrak{H}(C^*K)$ with a natural multiplicative structure, under the hypotheses of the proposition above.

2.3 The homotopy orbit model for free loop spaces.

Our goal here is to combine the thin free loop space model (2.2.4) with the general homotopy orbit space model of section 1.4, obtaining an extension $(\Lambda\upsilon \otimes C^*K \otimes \mathcal{B}C^*K, \widetilde{D})$ of $(\Lambda\upsilon, 0)$ by $fls^*(X)$, together with a quasi-isomorphism

$$\widetilde{\Upsilon} : (\Lambda\upsilon \otimes C^*K \otimes \mathcal{B}C^*K, \widetilde{D}) \longrightarrow (\Lambda\upsilon \otimes CU^*\mathcal{L}X, D^\sharp)$$

such that

$$\begin{array}{ccccc} (\Lambda v,0) & \xrightarrow{incl.} & (\Lambda v\otimes C^*K\otimes \mathcal{B}C^*K,\widetilde{D}) & \xrightarrow{\pi} & C^*K\odot \mathcal{B}C^*K \\ \| & & \downarrow{\scriptstyle \widetilde{\Upsilon}} & & \downarrow{\scriptstyle \Upsilon} \\ (\Lambda v,0) & \xrightarrow{incl.} & (\Lambda v\otimes CU^*\mathcal{L}X,D^\sharp) & \xrightarrow{\pi} & CU^*\mathcal{L}X \end{array} \tag{2.3.1}$$

commutes, where π denotes the obvious projections. We can construct the extension $(\Lambda v\otimes C^*K\otimes \mathcal{B}C^*K,\widetilde{D})$ under the additional hypothesis that the multiplication on C^*K is trivial. We show that the lift $\widetilde{\Upsilon}$ exists when K is such that $\psi_K^\sharp$ is commutative and the multiplication on C^*K is trivial and when the coefficient ring R is a field.

Restriction. Henceforth, we consider only those simplicial sets K such that $\psi_K^\sharp$ is commutative and such that C^*K has trivial multiplication. We suspect that our methods can be generalized to a larger class of spaces, though the formulas involved may be too complex to allow actual calculations to be performed. Furthermore, since the class of spaces we consider includes all simplicial suspensions $K = EL$ such that C_*L is cocommutative, as explained in the introduction to this section, we can actually treat a number of interesting spaces, including all double suspensions.

We now define the desired extension

$$(\Lambda v,0)\longrightarrow(\Lambda v\otimes C^*K\otimes \mathcal{B}C^*K,\widetilde{D})\longrightarrow C^*K\odot\mathcal{B}C^*K\ .$$

For any free graded module V, let $N:\perp V\longrightarrow\perp V$ denote the endomorphism given by

$$N(v_1|\cdots|v_n)=\sum_{1\le j\le n}\pm\, v_j|\cdots|v_n|v_1|\cdots|v_{j-1}$$

where the sign of each summand is determined by the Koszul rule, i.e., N associates to any word the signed sum of its cyclic permutations.

Proposition 2.3.1. *Let K be a 1-reduced simplicial set of finite type such that $\psi_K^\sharp$ is commutative and such that C^*K has trivial multiplication. Let S be the endomorphism of $C^*K\otimes\perp sC^+K$ of degree -1 given recursively for all $y\in C^+K$ and $sx_1|\cdots|sx_n\in\perp sC^+K$ by*

$$S(y\otimes 1)=1\otimes sy,$$

$$S(1\otimes sx_1|\cdots|sx_n)=0$$

and

$$\begin{aligned} S(y\otimes sx_1|\cdots|sx_n)=&1\otimes N(sy|sx_1|\cdots|sx_n)\\ &+\sum_{j\ge 1}S\big(\pi(sy\star sx_1\star\cdots\star sx_j)\otimes sx_{j+1}|\cdots|sx_n\big). \end{aligned}$$

Then $S^2=0$ and $\check{D}S=-S\check{D}$. Consequently,

$$\widetilde{D}=1\otimes\check{D}+v\otimes S$$

is a differential on $\Lambda v \otimes C^*K \otimes \mathcal{B}C^*K$.

Proof. An easy inductive proof establishes that $S^2 = 0$. Straightforward calculations then show that $\check{D}$ commutes with S. □

When K is a double suspension, it follows from the chain-level Bott-Samelson Theorem of [HPS2] that $\psi_K^\sharp$ is the shuffle product, which implies that $S(y \otimes sx_1|\cdots|sx_n) = 1 \otimes N(sy|sx_1|\cdots|sx_n)$.

To show that $(\Lambda v \otimes C^*K \otimes \mathcal{B}C^*K, \widetilde{D})$ has the desired cohomology, at least as a graded module, we prove that $(\Lambda v \otimes C^*K \otimes \mathcal{B}C^*K, \widetilde{D})$ is isomorphic as a cochain complex to the negative cyclic complex of the commutative algebra C^*K, looked at as a cochain complex in positive degrees, rather than as a chain complex in negative degrees. Jones proved in [J] that $\mathrm{H}^*(\mathcal{L}X_{hS^1})$ is isomorphic to the negative cyclic homology of the algebra S^*X, and therefore to that of C^*K, if $|K| \simeq X$. Thus, as a consequence of the next proposition, we obtain that

$$\mathrm{H}^*(\Lambda v \otimes C^*K \otimes \mathcal{B}C^*K, \widetilde{D}) \cong \mathrm{H}^*(\mathcal{L}X_{hS^1}),$$

as desired.

Recall that the negative cyclic complex on a cochain algebra (A, d), seen as a cochain complex, is

$$\mathfrak{C}(A) = (\Lambda v \otimes A \otimes \perp sA^+, D_{\mathfrak{C}}).$$

Here, v is, as always, of degree 2 and

$$D_{\mathfrak{C}} = 1 \otimes D_{\mathfrak{H}} + v \otimes S_{\mathfrak{C}},$$

where $S_{\mathfrak{C}}(b \otimes 1) = 1 \otimes sb$, $S_{\mathfrak{C}}(1 \otimes sa_1|\cdots|sa_n) = 0$, and

$$S_{\mathfrak{C}}(b \otimes sa_1|\cdots|sa_n) = 1 \otimes N(sb|sa_1|\cdots|sa_n).$$

Proposition 2.3.2. *Let* K *be a* 1*-reduced simplicial set of finite type such that* $\psi_K^\sharp$ *is commutative and such that* C^*K *has trivial multiplication. Let*

$$\mathfrak{h} : C^*K \odot \mathcal{B}C^*K \longrightarrow \mathfrak{H}(C^*K)$$

denote the isomorphism of Proposition 2.2.2. Let $\mathfrak{C}(C^*K)$ *denote the negative cyclic complex on* C^*K*, considered as a cochain complex. Then*

$$1 \otimes \mathfrak{h} : (\Lambda v \otimes C^*K \odot \mathcal{B}C^*K, \widetilde{D}) \longrightarrow \mathfrak{C}(C^*K)$$

is an isomorphism of cochain complexes.

Note that the two complexes are in fact equal when $\psi_K^\sharp$ is the shuffle product, e.g., when K is a double suspension.

Proof. Notice that

$$\begin{aligned}(1 \otimes \mathfrak{h})\widetilde{D} =& (1 \otimes \mathfrak{h})(1 \otimes \check{D} + v \otimes S)\\ =& 1 \otimes D_{\mathfrak{H}}\mathfrak{h} + v \otimes \mathfrak{h}S,\end{aligned}$$

while

$$D_{\mathfrak{C}}(1\otimes\mathfrak{h}) = 1\otimes D_{\mathfrak{H}}\mathfrak{H} + \upsilon\otimes S_{\mathfrak{C}}\mathfrak{h}.$$

To complete the proof, we need therefore to establish that $\mathfrak{h}S = S_{\mathfrak{C}}\mathfrak{h}$, which follows from an easy argument by induction on wordlength in $\perp sC^+K$. $\square$

Our principal reason to prefer $(\Lambda\upsilon\otimes C^*K\otimes\mathcal{B}C^*K,\widetilde{D})$ to $\mathfrak{C}(C^*K)$ concerns its application to further constructions. In [HR], the model $(\Lambda\upsilon\otimes C^*K\otimes\mathcal{B}C^*K,\widetilde{D})$ plays an essential role in the development of a model for calculating the spectrum homology of the topological cyclic homology of X, which requires the existence of a cochain quasi-isomorphism into $(\Lambda\upsilon\otimes CU^*\mathcal{L}X, D^\sharp)$ lifting a quasi-isomorphism into $CU^*(\mathcal{L}X)$. The model of the homotopy orbits used in [HR] must therefore lift the thin free loop space model of section 2.2, which $\mathfrak{C}(C^*K)$ does not.

Theorem 2.3.3. *Let K be a 1-reduced simplicial set of finite type such that the multiplication on C^*K is trivial and $\psi_K^\sharp$ is commutative. If the base ring R is a field, there is a quasi-isomorphism $\widetilde{\Upsilon}:(\Lambda\upsilon\otimes C^*K\otimes\mathcal{B}C^*K,\widetilde{D})\longrightarrow(\Lambda\upsilon\otimes CU^*\mathcal{L}X,D^\sharp)$ such that*

$$\begin{array}{ccccc}
(\Lambda\upsilon,0) & \xrightarrow{\text{incl.}} & (\Lambda\upsilon\otimes C^*K\otimes\mathcal{B}C^*K,\widetilde{D}) & \xrightarrow{\pi} & C^*K\odot\mathcal{B}C^*K \\
\| & & \downarrow{\scriptstyle\widetilde{\Upsilon}} & & \downarrow{\scriptstyle\Upsilon} \\
(\Lambda\upsilon,0) & \xrightarrow{\text{incl.}} & (\Lambda\upsilon\otimes CU^*\mathcal{L}|K|,D^\sharp) & \xrightarrow{\pi} & CU^*\mathcal{L}|K|
\end{array}$$

commutes, where π denotes the obvious projection maps.

Definition. Let X be a 1-connected space with the homotopy type of a finite-type CW-complex, and let K be a finite-type, 1-reduced simplicial set such that $|K|\simeq X$. Suppose that the natural product $\psi_K^\sharp$ on $\mathcal{B}C^*K$ is commutative and that the multiplication on C^*K is trivial. The twisted algebra extension

$$hos^*(X) := (\Lambda\upsilon\otimes C^*K\otimes\mathcal{B}C^*K,\widetilde{D})$$

of Theorem 2.3.3, together with the quasi-algebra quasi-isomorphism

$$\widetilde{\Upsilon}:(\Lambda\upsilon\otimes C^*K\otimes\mathcal{B}C^*K,\widetilde{D})\longrightarrow(\Lambda\upsilon\otimes CU^*\mathcal{L}X,D^\sharp),$$

is a *thin equivariant free loop model* for X.

The proof of Theorem 2.3.3 is based on the following result in homological perturbation theory, which we prove in the appendix.

Lemma 2.3.4. *Let $\Bbbk$ be any field. Let (C,d) and (C',d') be filtered cochain complexes of finite type over $\Bbbk$, where the filtrations $\{F_n\}_{n\geq 0}$ and $\{F'_n\}_{n\geq 0}$ are increasing and bounded below. Let $\varphi:(C,d)\to(C',d')$ be a filtration-preserving quasi-isomorphism such that the restriction $\varphi_n:F_n\to F'_n$ is a quasi-isomorphism for all n. Let $\theta:C\to C$ and $\theta':C'\to C'$ be filtration-decreasing linear maps of degree -1 such that $(d+\theta)^2=0$ and $(d'+\theta')^2=0$, i.e., $(C,d+\theta)$ and $(C',d'+\theta')$ are cochain complexes. Then there exists a filtration-decreasing linear map $\omega:C\to C'$ of degree 0 such that $\varphi+\omega:(C,d+\theta)\to(C',d'+\theta')$ is a quasi-isomorphism.*

Proof of Theorem 2.3.3. Consider the following differential filtrations, both of which are increasing and bounded below.

$$F_n\big((\Lambda\upsilon,0)\otimes(C^*K\odot\mathcal{B}C^*K)\big) := \Lambda\upsilon\otimes\big((C^*K\otimes\mathcal{B}C^*K)^{\leq n}\oplus(\operatorname{Im}\check{D})^{n+1}\big)$$

$$\widetilde{F}_n\big((\Lambda\upsilon,0)\otimes CU^*\mathcal{L}X\big) := \Lambda\upsilon\otimes\big((CU^{\leq n}\mathcal{L}X)\oplus(\operatorname{Im}d^\sharp)^{n+1}\big)$$

The cochain map $Id\otimes\Upsilon$ is clearly filtration-preserving with respect to these filtrations, and its restriction to each F_n is indeed a quasi-isomorphism. Furthermore, the differentials $\widetilde{D}$ and $D^\sharp$ are derived from the untwisted tensor-product differentials by filtration-decreasing perturbation. Lemma 2.3.4 therefore implies that there exists a filtration-decreasing linear map

$$\omega : \Lambda\upsilon\otimes C^*K\otimes\mathcal{B}C^*K \longrightarrow \Lambda\upsilon\otimes CU^*\mathcal{L}X$$

such that

$$Id\otimes\Upsilon+\omega : (\Lambda\upsilon\otimes C^*K\otimes\mathcal{B}C^*K,\widetilde{D}) \xrightarrow{\simeq} (\Lambda\upsilon\otimes CU^*\mathcal{L}X, D^\sharp)$$

is a quasi-isomorphism. We can therefore set $\widetilde{\Upsilon}=Id\otimes\Upsilon+\omega$. □

Related work. Bökstedt and Ottosen have recently developed an approach to Borel cohomologly calculations for free loop spaces that is Eckmann-Hilton dual to the approach considered here and thus complementary to our methods [BO]. They have constructed a Bousfield-type spectral sequence that converges to the cohomology of $(\mathcal{L}X)_{hS^1}$. While our model is easiest to deal with for spaces with few cells, the elementary cases for their model are Eilenberg-MacLane spaces.

2.4 Examples

We conclude this article with examples illustrating the use of the models we have built.

Example 2.4.1. Let $n>0$, and let K be the simplicial model of S^{2n+1} with exactly one nondegenerate simplex z of positive dimension, in dimension $2n+1$. We then have that $C^*K=\Lambda z$, with trivial differential. Furthermore, $\mathcal{B}C^*K$ is isomorphic as an algebra to Γsz, for degree reasons, and

$$fls^*(S^{2n+1}) = (\Lambda z\otimes\Gamma sz, 0).$$

Thus, $\mathrm{H}^*(\mathcal{L}S^{2n+1})\cong\Lambda z\otimes\Gamma sz$, as has long been known.

Let $sz(m)=sz|\cdots|sz\in\perp^m sz$. We then have

$$hos^*(S^{2n+1}) = (\Lambda\upsilon\otimes\Lambda z\otimes\Gamma sz,\widetilde{D})$$

where

$$\widetilde{D}(\upsilon^k\otimes 1\otimes sz(m))=0$$

for all k and m, while

$$\widetilde{D}(\upsilon^k\otimes z\otimes sz(m)) = (m+1)\upsilon^{k+1}\otimes 1\otimes sz(m+1).$$

The integral cohomology of the homotopy orbit space is therefore

$$\mathrm{H}^*(\mathcal{L}S^{2n+1}_{hS^1}) \cong \Lambda v \oplus \Gamma sz \oplus \bigoplus_{k,m\geq 1} \mathbb{Z}/m\mathbb{Z}\cdot(v^k\otimes sz(m))$$

as graded modules, while its mod p cohomology is of the form

$$\begin{aligned}\mathrm{H}^*(\mathcal{L}S^{2n+1}_{hS^1};\mathbb{F}_p) \cong &\Lambda v\otimes\mathbb{F}_p \oplus \Gamma sz\otimes\mathbb{F}_p\\ &\oplus \bigoplus_{k,m\geq 1}\mathbb{F}_p\cdot(v^k\otimes 1\otimes sz(pm))\\ &\oplus \bigoplus_{k,m\geq 1}\mathbb{F}_p\cdot(v^k\otimes z\otimes sz(pm-1))\end{aligned}$$

where a Bockstein sends the class of $v^k\otimes z\otimes sz(m)$ to the class of $v^{k+1}\otimes 1\otimes sz(m+1)$.

Example 2.4.2. Let K be the simplicial model of S^{2n} with exactly one nondegenerate simplex z of positive dimension, in dimension $2n$. We then have that $C^*K = \mathbb{Z}[z]/(z^2)$, with trivial differential. Furthermore, $\mathcal{B}C^*K$ is $\perp sz$, endowed with the shuffle product. Consequently,

$$fls^*(S^n) = (\mathbb{Z}[z]/(z^2)\otimes \perp sz, \check{D}),$$

where

$$\begin{aligned}\check{D}(1\otimes sz(m)) =& z\otimes sz(m-1) - (-1)^{m-1}z\otimes sz(m-1)\\ \check{D}(z\otimes sz(m)) =& 0\end{aligned}$$

for all m, if n is even, whence

$$\mathrm{H}^*(\mathcal{L}S^{2n};\mathbb{F}_2) = \mathbb{F}_2[z]/(z^2)\otimes \perp sz,$$

another well-known result.

The differential in the homotopy orbit model satisfies

$$\widetilde{D}(v^k\otimes 1\otimes sz(m)) = \begin{cases} 0 & : \quad m \text{ odd}\\ 2\cdot v^k\otimes z\otimes sz(m) & : \quad m \text{ even}\end{cases}$$

and

$$\widetilde{D}(v^k\otimes z\otimes sz(m)) = \begin{cases} 0 & : \quad m \text{ odd}\\ v^{k+1}\otimes 1\otimes sz(m+1) & : \quad m \text{ even.}\end{cases}$$

From these formulas, it follows easily that

$$\begin{aligned}\mathrm{H}^*(\mathcal{L}S^{2n};\mathbb{F}_2) \cong &\mathbb{F}_2[v]\oplus \perp sz\\ &\oplus \bigoplus_{k,m\geq 0}\mathbb{F}_2\cdot(v^{k+1}\otimes z\otimes sz(2m+1))\\ &\oplus \bigoplus_{k,m\geq 1}\mathbb{F}_2\cdot(v^k\otimes 1\otimes sz(2m)).\end{aligned}$$

Appendix. Proof of Lemma 2.3.4

Before beginning the proof itself, we provide a brief overview of the construction and use of mapping cones of cochain maps, as they play a critical role in the proof. Given a morphism of cochain complexes $f : (A, d) \to (A', d')$, its *mapping cone* is the cochain complex $C_f := (A' \oplus sA, D_f)$, where $D_f a' = d'a'$ and $D_f sa = f(a) - s(da)$. With the obvious conventions, we can write this differential in matrix notation as

$$D_f = \begin{pmatrix} d' & f \\ 0 & -sd \end{pmatrix}.$$

Note that if f is injective, then the natural surjective map

$$\mathfrak{q}_f : C_f \xrightarrow{\simeq} \operatorname{coker} f,$$

defined to be the quotient map on A' and 0 on sA, is a quasi-isomorphism.

Given two cochain maps $f : (A, d) \to (A', d')$ and $g : (B, d) \to (B'd')$, a cochain map $h : C_f \to C_g$ can be written in matrix notation as

$$h = \begin{pmatrix} h_{11} & h_{12} \\ h_{21} & h_{22} \end{pmatrix},$$

where $h_{11} : A' \to B'$, $h_{12} : sA \to B'$, $h_{21} : A' \to sB$ and $h_{22} : sA \to sB$.

It is an easy exercise to prove the following lemma.

Lemma A.1. *Let $h : C_f \to C_g$. If $h_{21} = 0$, then h_{11} and $s^{-1}h_{22}s : A \to A'$ are cochain maps and*

$$\begin{array}{ccc} A & \xrightarrow{f} & A' \\ {\scriptstyle s^{-1}h_{22}s}\downarrow & & \downarrow{\scriptstyle h_{11}} \\ B & \xrightarrow{g} & B' \end{array}$$

commutes up to cochain homotopy, given by $h_{12}s : A \to B'$.

The converse is equally trivial to establish.

Lemma A.2. *If*

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ {\scriptstyle \alpha}\downarrow & & \downarrow{\scriptstyle \beta} \\ A' & \xrightarrow{g} & B' \end{array}$$

commutes up to cochain homotopy, given by $\gamma : A \to B'$ of degree -1, then

$$h = \begin{pmatrix} \beta & \gamma s^{-1} \\ 0 & s\alpha s^{-1} \end{pmatrix}$$

is a cochain map from C_f to C_g. Moreover, if α and β are quasi-isomorphisms, so is h.

Proof of Lemma 2.3.4. The proof proceeds by induction on filtration degree.

The filtration of (C,d) is denoted

$$0 = F_0 \subset F_1 \subset ... \subset F_n \subset F_{n+1} \subset \cdots,$$

while the filtration of (C',d') is denoted

$$0 = F'_0 \subset F'_1 \subset ... \subset F'_n \subset F'_{n+1} \subset \cdots.$$

Since the perturbations θ and θ' are filtration-lowering, the filtrations of (C,d) and of $(C'd')$ give rise to filtrations of $(C,d+\theta)$ and of $(C',d'+\theta')$, denoted by

$$0 = \widetilde{F}_0 \subset \widetilde{F}_1 \subset ... \subset \widetilde{F}_n \subset \widetilde{F}_{n+1} \subset \cdots$$

and

$$0 = \widetilde{F}'_0 \subset \widetilde{F}'_1 \subset ... \subset \widetilde{F}'_n \subset \widetilde{F}'_{n+1} \subset \cdots,$$

respectively.

Recall that we assume that $\varphi_n : F_n \xrightarrow{\simeq} F'_n$, the restriction of the quasi-isomorphism $\varphi : (C,d) \to (C',d')$ to F_n, is itself also a quasi-isomorphism, for all n. There is therefore a commuting diagram of short exact sequences of cochain complexes

$$\begin{array}{ccccccccc}
0 & \longrightarrow & F_n & \xrightarrow{\iota_n} & F_{n+1} & \xrightarrow{\pi_{n+1}} & F_{n+1}/F_n & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle \varphi_n}\simeq & & \downarrow{\scriptstyle \varphi_{n+1}}\simeq & & \downarrow{\scriptstyle \phi_{n+1}} & & \\
0 & \longrightarrow & F'_n & \xrightarrow{\iota'_n} & F'_{n+1} & \xrightarrow{\pi'_{n+1}} & F'_{n+1}/F'_n & \longrightarrow & 0
\end{array}$$

and so the induced map on filtration quotients $\phi_{n+1} : F_{n+1}/F_n \to F'_{n+1}/F'_n$ is also a quasi-isomorphism.

Observe that $\widetilde{F}_1 = F_1$ and $\widetilde{F}'_1 = F'_1$. Consequently, if we define $\widetilde{\varphi}_1 : \widetilde{F}_1 \to \widetilde{F}'_1$ to be φ_1, then $\widetilde{\varphi}_1$ is a quasi-isomorphism.

Suppose that for some $n \geq 2$, there is a quasi-isomorphism $\widetilde{\varphi}_{n-1} : \widetilde{F}_{n-1} \to \widetilde{F}'_{n-1}$ such that $\omega_{n-1} = \widetilde{\varphi}_{n-1} - \varphi_{n-1} \in \widetilde{F}'_{n-2}$. Consider the following commuting diagram of short exact sequences of cochain complexes.

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \widetilde{F}_{n-1} & \xrightarrow{\iota_{n-1}} & \widetilde{F}_n & \xrightarrow{\pi_n} & \widetilde{F}_n/\widetilde{F}_{n-1} = F_n/F_{n-1} & \longrightarrow & 0 \\
\downarrow & & \downarrow{\scriptstyle \widetilde{\varphi}_{n-1}}\simeq & & & & \downarrow{\scriptstyle \phi_n}\simeq & & \downarrow \\
0 & \longrightarrow & \widetilde{F}'_{n-1} & \xrightarrow{\iota'_{n-1}} & \widetilde{F}'_n & \xrightarrow{\pi'_n} & \widetilde{F}'_n/\widetilde{F}'_{n-1} = F'_n/F'_{n-1} & \longrightarrow & 0
\end{array}$$

It induces a commuting diagram of long exact sequences in cohomology

$$\begin{array}{ccccccccccccc}
\cdots & \longrightarrow & \mathrm{H}^{k-1}\, F_n/F_{n-1} & \xrightarrow{\partial_{k-1}} & \mathrm{H}^k\, \widetilde{F}_{n-1} & \xrightarrow{\mathrm{H}^k\, \iota_{n-1}} & \mathrm{H}^k\, \widetilde{F}_n & \xrightarrow{\mathrm{H}^k\, \pi_n} & \mathrm{H}^k\, F_n/F_{n-1} & \xrightarrow{\partial_k} & \mathrm{H}^{k+1}\, \widetilde{F}_{n-1} & \longrightarrow & \cdots \\
 & & \downarrow\cong & & \downarrow\cong & & & & \downarrow\cong & & \downarrow\cong & & \\
\cdots & \longrightarrow & \mathrm{H}^{k-1}\, F'_n/F'_{n-1} & \xrightarrow{\partial'_{k-1}} & \mathrm{H}^k\, \widetilde{F}'_{n-1} & \xrightarrow{\mathrm{H}^k\, \iota'_{n-1}} & \mathrm{H}^k\, \widetilde{F}'_n & \xrightarrow{\mathrm{H}^k\, \pi'_n} & \mathrm{H}^k\, F'_n/F'_{n-1} & \xrightarrow{\partial_k} & \mathrm{H}^{k+1}\, \widetilde{F}'_{n-1} & \longrightarrow & \cdots,
\end{array}$$

which in turn gives rise to a commuting diagram of short exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \operatorname{coker}\partial_{k-1} & \longrightarrow & \mathrm{H}^k\,\widetilde{F}_n & \longrightarrow & \ker\partial_k & \longrightarrow & 0 \\ & & \downarrow\cong & & & & \downarrow\cong & & \\ 0 & \longrightarrow & \operatorname{coker}\partial'_{k-1} & \longrightarrow & \mathrm{H}^k\,\widetilde{F}'_n & \longrightarrow & \ker\partial'_k & \longrightarrow & 0. \end{array}$$

Since we are working over a field, both short exact sequences split, so there is an isomorphism $\Phi_n^k : \mathrm{H}^k\,\widetilde{F}_n \xrightarrow{\cong} \mathrm{H}^k\,\widetilde{F}'_n$ that fits into the commuting diagram of long exact sequences.

Next choose any section of the surjection from the cocycles of $\widetilde{F}_{n-1}$ onto its cohomology, then postcompose with the inclusion of the cocycles into $\widetilde{F}_{n-1}$, to obtain a injective quasi-isomorphism

$$j_{n-1} : \mathrm{H}^*\,\widetilde{F}_{n-1} \xrightarrow{\simeq} \widetilde{F}_{n-1}.$$

Let

$$j'_{n-1} := \widetilde{\varphi}_{n-1} j_{n-1} (\mathrm{H}^*\,\widetilde{\varphi}_{n-1})^{-1} : \mathrm{H}^*\,\widetilde{F}'_{n-1} \xrightarrow{\simeq} \widetilde{F}'_{n-1},$$

so that

$$\begin{array}{ccc} \mathrm{H}^*\,\widetilde{F}_{n-1} & \xrightarrow[\simeq]{j_{n-1}} & \widetilde{F}_{n-1} \\ {\scriptstyle \mathrm{H}^*\,\widetilde{\varphi}_{n-1}}\downarrow\cong & & \widetilde{\varphi}_{n-1}\downarrow\simeq \\ \mathrm{H}^*\,\widetilde{F}'_{n-1} & \xrightarrow[\simeq]{j'_{n-1}} & \widetilde{F}'_{n-1} \end{array}$$

commutes. Define $k_n : \mathrm{H}^*\,F_n/F_{n-1} \xrightarrow{\simeq} F_n/F_{n-1}$ and $k'_n : \mathrm{H}^*\,F'_n/F'_{n-1} \xrightarrow{\simeq} F'_n/F'_{n-1}$ similarly, so that

$$\begin{array}{ccc} \mathrm{H}^*\,F_n/F_{n-1} & \xrightarrow[\simeq]{k_n} & F_n/F_{n-1} \\ {\scriptstyle \mathrm{H}^*\,\phi_n}\downarrow\cong & & \phi_n\downarrow\simeq \\ \mathrm{H}^*\,F'_n/F'_{n-1} & \xrightarrow[\simeq]{k'_n} & F'_n/F'_{n-1} \end{array}$$

commutes.

For each $s \geq 1$, fix a basis $\mathfrak{B}_s = \big([z_{s1}], ..., [z_{sm_s}]\big)$ of $\mathrm{H}^s\,\widetilde{F}_n$, where $[z_{si}]$ denotes the cohomology class of a cocycle $z_{si} \in \widetilde{F}_n^s$. Then there exists $w \in (F_n/F_{n-1})^{s-1}$ such that

$$k_n\,\mathrm{H}^s\,\pi_n\big([z_{si}]\big) = \pi_n(z_{si}) - \bar{d}w,$$

where $\bar{d}$ denotes the differential on F_n/F_{n-1}. Since π_n is surjective, there exists $v \in \widetilde{F}_n^{s-1}$ such that $\pi_n(v) = w$. Thus, if we set $j_n\big([z_{si}]\big) = z_{si} - (d+\theta)v$, we obtain that

$$\pi_n j_n\big([z_{si}]\big) = k_n\,\mathrm{H}^s\,\pi_n\big([z_{si}]\big).$$

Proceeding in this manner, we can define a quasi-isomorphism $j_n : H^*\widetilde{F}_n \xrightarrow{\simeq} \widetilde{F}_n$ such that

$$\begin{array}{ccc} \mathrm{H}^*\,\widetilde{F}_n & \xrightarrow[\simeq]{j_n} & \widetilde{F}_n \\ {\scriptstyle H^*\pi_n}\downarrow & & \pi_n\downarrow \\ H^*F_n/F_{n-1} & \xrightarrow[\simeq]{k_n} & F_n/F_{n-1} \end{array}$$

commutes. Similarly, we can construct a quasi-isomorphism $j'_n : H^*\widetilde{F}'_n \xrightarrow{\simeq} \widetilde{F}'_n$ such that

$$\begin{array}{ccc} \mathrm{H}^*\,\widetilde{F}'_n & \xrightarrow[\simeq]{j'_n} & \widetilde{F}'_n \\ {\scriptstyle H^*\pi'_n}\downarrow & & \downarrow{\scriptstyle \pi'_n} \\ H^*F'_n/F'_{n-1} & \xrightarrow[\simeq]{k'_n} & F'_n/F'_{n-1} \end{array}$$

commutes.

Observe that


$$\begin{array}{ccc} \mathrm{H}^*\,\widetilde{F}_{n-1} & \xrightarrow[\simeq]{j_{n-1}} & \widetilde{F}_{n-1} \\ {\scriptstyle H^*\iota_{n-1}}\downarrow & & \downarrow{\scriptstyle \iota_{n-1}} \\ \mathrm{H}^*\,\widetilde{F}_n & \xrightarrow[\simeq]{j_n} & \widetilde{F}_n \end{array}$$

and

$$\begin{array}{ccc} \mathrm{H}^*\,\widetilde{F}'_{n-1} & \xrightarrow[\simeq]{j'_{n-1}} & \widetilde{F}'_{n-1} \\ {\scriptstyle H^*\iota'_{n-1}}\downarrow & & \downarrow{\scriptstyle \iota'_{n-1}} \\ \mathrm{H}^*\,\widetilde{F}'_n & \xrightarrow[\simeq]{j'_n} & \widetilde{F}'_n \end{array}$$

necessarily commute up to cochain homotopy.

Uniting all of these elements, we have a diagram



in which all squares commute exactly, except for the top left and the bottom left squares, which commute up to cochain homotopy. We must now show that the missing arrow can be filled in, so that the two front squares thus obtained commute, at least up to cochain homotopy.


Since the top and bottom left squares commute up to homotopy, by Lemma A.2 there are induced quasi-isomorphisms

$$\mathfrak{j}_n : C_{\mathrm{H}^*\,\iota_{n-1}} \xrightarrow{\simeq} C_{\iota_{n-1}} \quad \text{and} \quad \mathfrak{j}_n : C_{\mathrm{H}^*\,\iota_{n-1}} \xrightarrow{\simeq} C_{\iota_{n-1}},$$


which, together with the induced isomorphism

$$\mathfrak{f}_n : C_{\mathrm{H}^*\,\iota_{n-1}} \xrightarrow{\cong} C_{\mathrm{H}^*\,\iota'_{n-1}},$$

give rise to a commutative diagram




Let $\mathfrak{p}_n : C_{\iota_{n-1}} \longrightarrow s\widetilde{F}_{n-1}$ and $\mathfrak{p}'_n : C_{\iota'_{n-1}} \longrightarrow s\widetilde{F}'_{n-1}$ denote the projection maps. The commutativity of the previous diagram implies that the following diagram commutes as well.



Since $\mathfrak{j}_n$ is an injective quasi-isomorphism and $\mathfrak{q}_{\iota'_{n-1}} + \mathfrak{p}'_n$ is a surjection, there is a cochain map

$$\mathfrak{g}_n : C_{\iota_{n-1}} \longrightarrow C_{\iota'_{n-1}}$$

such that



commutes. The lift $\mathfrak{g}_n$ must be a quasi-isomorphism because the upper triangle commutes. Writing $\mathfrak{g}_n$ in the matrix notation introduced at the beginning of this appendix as

$$\begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix},$$

we see that $g_{22} = s\widetilde{\varphi}_{n-1}s^{-1}$, since $\mathfrak{p}'_n\mathfrak{g}_n = s\widetilde{\varphi}_{n-1}s^{-1}\mathfrak{p}_n$. Furthermore the equality $\mathfrak{q}_{\iota'_{n-1}}\mathfrak{g}_n = \phi_n\mathfrak{q}_{\iota_n}$ implies that $g_{11} - \varphi_n \in \widetilde{F}'_{n-1}$.

It remains to show that $g_{21}=0$, as Lemma A.1 then implies that $g_{11}:\widetilde{F}_n\longrightarrow\widetilde{F}'_n$ is a chain map such that

$$\begin{array}{ccc}\widetilde{F}_{n-1} & \xrightarrow{\iota_{n-1}} & \widetilde{F}_n\\ {\scriptstyle\widetilde{\varphi}_{n-1}}\downarrow\simeq & & \downarrow{\scriptstyle g_{11}}\\ \widetilde{F}'_{n-1} & \xrightarrow{\iota'_{n-1}} & \widetilde{F}'_n\end{array}$$

commutes up to cochain homotopy, given by $g_{12}s:\widetilde{F}_{n-1}\longrightarrow\widetilde{F}'_n$. Since $\widetilde{\varphi}_{n-1}$ and $\mathfrak{g}_n$ are both quasi-isomorphisms, it follows then that g_{11} is also a quasi-isomorphism. Letting $\widetilde{\varphi}_n=g_{11}$, we complete the inductive step of the proof.

To see that $g_{21}=0$, note that, since we are working over a field, $C_{\iota_{n-1}}$ can be split as a cochain complex into $C_{\mathrm{H}^*\iota_{n-1}}\oplus W\oplus D_{\mathrm{H}^*\iota_{n-1}}W$. Given $w=x+sy\in W$, where $x\in\widetilde{F}_n$ and $y\in\widetilde{F}_{n-1}$, observe that

$$(\phi_n\oplus s\widetilde{\varphi}_{n-1}s^{-1})(\mathfrak{q}_{\iota_{n-1}}+\mathfrak{p}_n)(w)=\overline{\varphi_n(x)}+s\widetilde{\varphi}_{n-1}(y)=(\mathfrak{q}_{\iota'_{n-1}}+\mathfrak{p}'_n)\big(\varphi_n(x)+s\widetilde{\varphi}_{n-1}(y)\big),$$

where $\overline{\varphi_n(x)}$ denotes the class of $\varphi_n(x)$ in F'_n/F'_{n-1}. We can therefore set $\mathfrak{g}_n(w)=\varphi_n(x)+s\widetilde{\varphi}_{n-1}(y)$, which corresponds to $g_{11}(x)=\varphi_n(x)$, $g_{12}(sy)=0$, $g_{21}(x)=0$ and $g_{22}(sy)=(s\widetilde{\varphi}_{n-1}s^{-1})(sy)$. For $\mathfrak{g}_n$ to be a cochain map, it is now necessary and sufficient to define $\mathfrak{g}_n\big(D_{\mathrm{H}^*\iota_{n-1}}w\big)$ to be $D_{\iota'_{n-1}}\mathfrak{g}_n(w)$ for all $w\in W$. More explicitly,

$$\mathfrak{g}_n\big(D_{\mathrm{H}^*\iota_{n-1}}(x+sy)\big)=d\varphi_n(x)+\iota_{n-1}\widetilde{\varphi}_{n-1}(y)-s\big(d\widetilde{\varphi}_{n-1}(y)\big),$$

and thus $g_{21}=0$. □

References

[A] Anick, D., *Hopf algebras up to homotopy*, J. Amer. Math. Soc. **2** (1989), 417-453.

[B] Baues, H.-J., *The cobar construction as a Hopf algebra*, Invent. Math. **132** (1998), 467–489.

[Bl] Blanc, S., *Modèles tordus d'espaces de lacets libres et fonctionnels*, Thesis, EPFL (2004).

[BO] Bökstedt, M.; Ottosen, I., *A spectral sequence for string cohomology*, Topology **44** (2005), 1181–1212.

[DH] N. Dupont and K. Hess, *Commutative free loop space models at large primes*, Math. Z. **244** (2003), 1-34.

[FHT] Félix, Y.; Halperin, S.; Thomas, J.-C., *Differential graded algebras in topology*, Handbook of Algebraic Topology, Elsevier, 1995, pp. 829-865.

[G] Getzler, E., *Batalin-Vilkovisky algebras and two-dimensional topological field theories*, Comm. Math. Phys. **159** (1994), 265–285.

[H] Hess, K., *An algebraic model for mod 2 topological cyclic homology*, String Topology and Cyclic Homology, Birkhaüser, 2006, pp. 97–162.

[HPS1] Hess, K.; Parent, P.-E.; Scott, J., *Bimodules over operads characterize morphisms*, preprint at arXiv:math.AT/0505559 (2005).

[HPS2] Hess, K.; Parent, P.-E.; Scott, J., *The canonical Adams-Hilton model of a suspension*, In preparation.

[HPST] Hess, K.; Parent, P.-E.; Scott, J.; Tonks, A., *A canonical enriched Adams-Hilton model for simplicial sets*, To appear in *Adv. in Math.*, preprint at arXiv:math.AT/0408216 (2004).

[HR] Hess, K.; Rognes, J., *Algebraic models for topological cyclic homology and Whitehead spectra of simply connected spaces*, In preparation.

[K] Kontsevich, M., *Operads and motives in deformation quantization*, Lett. Math. Phys. **48** (1999), 35–72.

[Mas] Massey, W., *Singular Homology Theory*, Graduate Texts in Mathematics, vol. 70, Springer, 1980.
[M] May, J.P., *Simplicial Objects in Algebraic Topology*, Midway reprint 1982, University of Chicago Press, 1967.
[Mi] Milgram, R.J., *Iterated loop spaces*, Ann. of Math. **84** (1966), 386–403.
[Mo] Moore, J.C., *Algèbre homologique et homologie des espaces classifiants*, Exposé 7, Séminaire Cartan 1959/60.
[S] Szczarba, R.H., *The homology of twisted cartesian products*, Trans. Amer. Math. Soc. **100** (1961), 197–216.

Institut de géométrie, algèbre et topologie (IGAT), Ecole Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland
E-mail address: kathryn.hess@epfl.ch